



# A strictly contractive Peaceman-Rachford splitting method for the doubly nonnegative relaxation of the minimum cut problem

Xinxin Li<sup>1</sup> · Ting Kei Pong<sup>2</sup> · Hao Sun<sup>3</sup> · Henry Wolkowicz<sup>3</sup> 

Received: 22 October 2019 / Accepted: 24 December 2020

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC part of Springer Nature 2021

## Abstract

The minimum cut problem, **MC**, and the special case of the vertex separator problem, consists in partitioning the set of nodes of a graph  $G$  into  $k$  subsets of given sizes in order to minimize the number of edges cut after removing the  $k$ -th set. Previous work on approximate solutions uses, in increasing strength and expense: eigenvalue, semidefinite programming, **SDP**, and doubly nonnegative, **DNN**, bounding techniques. In this paper, we derive strengthened **SDP** and **DNN** relaxations, and we propose a scalable algorithmic approach for efficiently evaluating, theoretically verifiable, both upper and lower bounds. Our stronger relaxations are based on a new *gangster set*, and we demonstrate how *facial reduction*, **FR**, fits in well to allow for *regularized* relaxations. Moreover, the **FR** appears to be perfectly well suited for a *natural* splitting of variables, and thus for the application of *splitting* methods. Here, we adopt the strictly contractive Peaceman-Rachford splitting method, **sPRSM**. Further, we bring *useful* redundant constraints back into the subproblems, and show empirically that this accelerates **sPRSM**. In addition, we employ new strategies for obtaining lower bounds and upper bounds of the optimal value of **MC** from approximate iterates of the **sPRSM** thus aiding in early termination of the algorithm. We compare our approach with others in the literature on random datasets and vertex separator problems. This illustrates the efficiency and robustness of our proposed method.

**Keywords** Semidefinite relaxation · Doubly nonnegative relaxation · Min-cut · Graph partitioning · Vertex separator · Peaceman-Rachford splitting method · Facial reduction

---

This paper is partially based on the Master's thesis of Hao Sun [27]. The authors Hao Sun and Henry Wolkowicz thank the Natural Sciences and Engineering Research Council of Canada for their support. Xinxin Li's research was supported by the National Natural Science Foundation of China (No. 11601183, No. 61872162), Natural Science Foundation for Young Scientist of Jilin Province (No. 20180520212JH) and the China Scholarship Council (No. 201806175127). Ting Kei Pong's research was supported partly by Hong Kong Research Grants Council PolyU153004/18p.

---

Extended author information available on the last page of the article

## 1 Introduction

We present strengthened doubly nonnegative, **DNN**, (both positive semidefinite, **SDP**, and nonnegative elementwise) relaxations for the min-cut problem, **MC**; i.e., for the problem of partitioning the set of nodes of a graph  $G$  into  $k$  subsets of given sizes in order to minimize the number of edges cut after removing the  $k$ -th set. Our relaxations are aimed at specifically applying splitting methods based on using the regularization technique facial reduction, **FR**. We see that the **FR** technique both provides an excellent splitting of variables, as well as reducing the singularity degree to zero (equivalent to strict feasibility) and thus providing regularization for improving accuracy. In addition, we employ new so-called *gangster constraints*. The result is strengthened and theoretically verifiable upper and lower bounds for **MC**.

We consider an *undirected graph*  $G = (\mathcal{V}, \mathcal{E})$  with vertex and edge sets  $\mathcal{V}$ ,  $\mathcal{E}$ , respectively, and  $|\mathcal{V}| = n$ . We let  $m = (m_1 \ m_2 \ \dots \ m_k)^T$ ,  $\sum_{i=1}^k m_i = n$ , denote a given partition of  $n$  into  $k$  sets. The special type of *minimum cut problem*, **MC**, we consider consists in partitioning the vertex set  $\mathcal{V}$  into  $k$  subsets, with given sizes in  $m$ , in order to *minimize the cut* obtained after removing the  $k$ -th set, i.e., we minimize the number of edges connecting distinct sets other than those edges connected to the  $k$ -th set, see e.g., [21]. This problem can be modelled as a linearly constrained quadratic 0, 1 program

$$\begin{aligned} \text{cut}(m) = \min & \quad \frac{1}{2} \text{trace} AXBX^T \\ \text{s.t.} & \quad X \in \mathcal{M}_m, \end{aligned} \quad (1.1)$$

where the matrices  $A$ ,  $B$  and the set of 0, 1 partition matrices  $\mathcal{M}$  are defined below in Sect. 1.2, see (1.2).

This problem arises for example when finding an *ordering* to bring the sparsity pattern of a large sparse positive definite matrix into a block-arrow shape so as to minimize fill-in within a Cholesky factorization, e.g., [4, 9, 26]. The **MC** has further applications in computer program segmentation, solving symmetric systems of equations, microchip design and circuit board, floor planning and other layout problems [20]. In particular herein, we include consideration of the vertex separator problem, i.e., finding a vertex set whose removal splits the graph into *two* disconnected subsets, see e.g., [8, 22].

It is well known that **MC** is an NP-hard problem when  $k \geq 3$ , see e.g., [15, 21]. Solution techniques rely on efficiently calculating lower and upper bounds. We refer the readers to [7, 10, 19, 21, 22] and the references therein for recent results for finding bounds and solving **MC**; and also to [22, Sect. 2] for a recent overview of existing relaxation techniques for solving **MC**. An important tool for finding lower bounds is the *semidefinite programming*, **SDP**, relaxation of **MC**; this is included in [19]. Moreover, this relaxation uses *facial reduction* **FR** to guarantee strict feasibility and robustness for both the relaxation and its dual. However, these **SDP** problems are typically solved by interior point methods. These methods often do not scale

well, cannot properly exploit sparsity, and generally do *not* provide high accuracy solutions. Moreover, while lower bounds from **SDP** can be strengthened to yield better approximations to **MC** by adding extra nonnegativity and cutting plane constraints, the resulting optimization problems can be prohibitively expensive to solve for interior point solvers. And, the lack of accuracy in solutions means that both lower and upper bounds are *not* necessarily true bounds. Thus, in order to improve **MC** approximations, besides deriving tighter theoretical upper and lower bounds, one also needs to design efficient, accurate, and scalable algorithms for computing these bounds.

## 1.1 Main contributions

In this paper, we derive tighter (lower and upper) bounds and design efficient algorithms for their evaluation. The bounds are based on strengthened **SDP** and doubly nonnegative, **DNN**, relaxations within a **FR** framework. The bounding techniques are theoretically verifiable. Moreover, we introduce a random weighted sampling of eigenvectors to strengthen the upper bounds.

Our stronger relaxations use a new *gangster* set; see Definition 2.3. This set can be larger than the one used in the literature, e.g., [19, 29], when some of the set sizes  $m_i = 1$ . Then we show that the facially reduced problems satisfy the Robinson regularity condition. In addition, we show that many of the constraints are redundant in the facially reduced problem.

Our final **DNN** relaxation can be very difficult and time consuming to solve for interior point solvers. Therefore, we propose a scalable and regularized algorithmic approach. The key idea is that **FR** gives a natural way of reformulating the facially reduced **DNN** relaxation into a *separable* convex programming problem with linear coupling constraints. This sets the stage for an application of *splitting methods* such as *alternating direction method of multipliers*, **ADMM** [3]. These methods typically involve updating the multiplier(s) and solving several subproblems at each iteration. Their efficiency depends highly on the simplicity of the subproblems, and accurate solutions can take many iterations.

Herein we employ a particular variant of **ADMM**, the *strictly contractive Peaceman-Rachford splitting method*, **sPRSM**, [11, 12]. This method involves two subproblems and two updates of the multiplier at every iteration. While a direct application of this method can be slow (i.e., takes a lot of iterations), we introduce *two* key ingredients for empirical acceleration. First, instead of just using the natural splitting induced by **FR**, as in the recent work [18], we bring back some provably redundant constraints that are *not* redundant for the subproblems and do not significantly increase the computational cost. Second, we derive new strategies for obtaining lower and upper bounds of the true optimal value of **MC**. This helps with early termination of **sPRSM** when the two bounds agree. Specifically, we compute a lower bound by looking at the Fenchel dual. Moreover, we mimic the now classical Goeman-Williamson's approach for MAXCUT and use a random weighted sampling of eigenvectors of an iterate of the **sPRSM** before projecting it onto the set of partition matrices for computing an upper bound.

In the numerical experiments, we illustrate the efficiency of our proposed algorithmic approach (based on the strengthened **DNN** relaxation model) by comparing with the **DNN** relaxation model in [19], as well as the **SDP**<sub>4</sub> model in [22]. Our experiments show that our approach takes less computational time and the bounds obtained are generally strengthened.

### 1.1.1 Outline

In Sect. 2 we discuss properties of our new gangster sets and our facially reduced **SDP** and **DNN** relaxations. Our algorithmic **sPRSM** approach is presented in Sect. 3. In addition, we discuss the usefulness of redundant constraints and include details of the subproblems of **sPRSM**. And, we describe the methods for obtaining both lower and upper bounds from possibly inaccurate solutions of the **sPRSM**. Our numerical results are presented in Sect. 4. Concluding remarks are given in Sect. 5.

## 1.2 Preliminaries

Let  $A$  be the adjacency matrix of our graph,  $G = (\mathcal{V}, \mathcal{E})$ . Let  $e_j, E_j = e_j e_j^T, I_j$  denote, respectively, the all ones vector (of dimension  $j$ ), all ones matrix, and identity matrix. We use  $e, E, I$ , when the dimensions are clear. We set

$$B = \begin{bmatrix} E_{k-1} & -I_{k-1} & 0 \\ 0 & 0 & 0 \end{bmatrix} \in \mathbb{S}^k,$$

where  $\mathbb{S}^k$  is the space of real symmetric  $k \times k$  matrices equipped with the *trace inner product*,  $\langle S, T \rangle = \text{trace } ST$ , and the corresponding *Fröbenius norm*,  $\|S\|_F$ . We use  $\|S\| = \|S\|_F$ , when the meaning is clear.

Let  $m = (m_1, \dots, m_k)^T \in \mathbb{Z}_+^k, k > 2$ , and let  $n = |\mathcal{V}| = m^T e$ . Let  $S = \{S_1, S_2, \dots, S_k\}$  be a partition of the vertex set with cardinalities  $|S_i| = m_i > 0, i = 1, \dots, k$ , i.e., the sets are nonempty, pairwise disjoint, and the union is  $S$ . In addition, we let  $M = \text{Diag}(m)$  denote the diagonal matrix formed from the vector  $m$ . More generally, for a vector  $x \in \mathbb{R}^j$ , we define  $\text{Diag} : \mathbb{R}^j \rightarrow \mathbb{S}^j$  to be the linear transformation that maps  $x$  to the diagonal matrix whose diagonal is  $x$ ; we denote its adjoint linear transformation by  $\text{diag}$ , i.e.  $\text{diag} := \text{Diag}^*$ . Next, we define the set of edges between two sets of nodes by

$$\delta(S_i, S_j) := \{uv \in \mathcal{E} : u \in S_i, v \in S_j\}.$$

The *cut of a partition*  $S$ ,  $\delta(S)$ , is then defined as the union of all edges cut by the first  $k - 1$  sets of the partition, i.e.,

$$\delta(S) := \cup \{ \delta(S_i, S_j) : 1 \leq i < j \leq k - 1 \}.$$

Our objective is to minimize the cardinality of the cut, i.e.,  $|\delta(S)|$ . In [21], it is shown that  $|\delta(S)|$  can be represented in terms of a quadratic form of the partition matrix  $X$ . This quadratic form for the **MC** problem in the trace formulation is

$$\begin{aligned} \text{cut}(m) &= \min \frac{1}{2} \text{trace} AXBX^T \\ \text{s.t.} \quad & X \in \mathcal{M}_m, \end{aligned} \tag{1.2}$$

where the set of *partition matrices*,  $\mathcal{M}_m$  is defined by

$$\mathcal{M}_m = \{X \in \mathbb{R}^{n \times k} : Xe = e, X^T e = m, X_{ij} \in \{0, 1\}\},$$

i.e., column  $j$  of a partition matrix  $X$  is the *indicator vector* for set  $S_j$ . We let  $x = \text{vec}(X) \in \mathbb{R}^{nk}$  denote the columnwise vectorization of the matrix  $X$ . The inverse and *adjoint linear transformation*  $\text{Mat} : \mathbb{R}^{nk} \rightarrow \mathbb{R}^{n \times k}$  is

$$X = \text{Mat}(x) = \text{vec}^*(x) = \text{vec}^{-1}(x).$$

One way to derive an **SDP** relaxation for (1.2) is to start by considering a Lagrangian relaxation of a quadratic-quadratic model of **MC**. Taking the dual of the dual of this Lagrangian relaxation then gives the **SDP** relaxation for (1.2); see also [29, 31] for the development for other hard combinatorial problems. Alternatively, we can obtain the *same SDP* relaxation directly using the well-known *lifting process*, e.g., [2, 16, 25, 29, 31].

We denote the matrix lifting

$$Y := \begin{pmatrix} 1 \\ x \end{pmatrix} \begin{pmatrix} 1 \\ x \end{pmatrix}^T, \quad x = \text{vec}(X). \tag{1.3}$$

Then  $Y \in \mathbb{S}_+^{nk+1}$  and is rank-one. The rows and columns of  $Y$  are indexed from 0 to  $nk$ . Note that  $Y$  in (1.3) can be blocked appropriately as

$$Y = \begin{bmatrix} Y_{00} & Y_{1:nk 0}^T \\ Y_{1:nk 0} & \bar{Y} \end{bmatrix}, \quad Y_{1:nk 0} = \begin{bmatrix} Y_{(10)} \\ Y_{(20)} \\ \vdots \\ Y_{(k0)} \end{bmatrix}, \quad \bar{Y} = \begin{bmatrix} \bar{Y}_{(11)} & \bar{Y}_{(12)} & \cdots & \bar{Y}_{(1k)} \\ \bar{Y}_{(21)} & \bar{Y}_{(22)} & \cdots & \bar{Y}_{(2k)} \\ \vdots & \ddots & \ddots & \vdots \\ \bar{Y}_{(k1)} & \ddots & \ddots & \bar{Y}_{(kk)} \end{bmatrix}, \tag{1.4}$$

with

$$\bar{Y}_{(ij)} \in \mathbb{R}^{n \times n}, \forall i \neq 0, \forall j \neq 0, \text{ and } Y_{(j0)} \in \mathbb{R}^n, \forall j = 1, \dots, k.$$

With the matrix lifting for  $Y$ , we rewrite the objective function in (1.2) in the linearized form:

$$\frac{1}{2} \text{trace} AXBX^T = \frac{1}{2} \text{trace} L_A Y, \quad L_A := \begin{bmatrix} 0 & 0 \\ 0 & B \otimes A \end{bmatrix}. \tag{1.5}$$

In [19, 29], a preliminary **SDP** relaxation is derived for the model (1.2). After applying facial reduction to the **SDP** relaxations, the variable  $Y$  is expressed as  $Y = \tilde{V}R\tilde{V}^T$ , for some full column rank matrix  $\tilde{V} \in \mathbb{R}^{(nk+1) \times ((k-1)(n-1)+1)}$ , and the following **SDP** is obtained in [19] after removing some redundant constraints:

$$\begin{aligned} \min \quad & \frac{1}{2} \text{trace} \left( \tilde{V}^T L_A \tilde{V} \right) R \\ \text{s.t.} \quad & \mathcal{G}_{\hat{J}}(\tilde{V} R \tilde{V}^T) = u_0 \\ & R \geq 0, \end{aligned} \tag{1.6}$$

where the linear transformation  $\mathcal{G}_{\hat{J}}(\cdot)$  is called the gangster operator and  $u_0$  is the first unit vector. The Slater constraint qualification, holds for both (1.6) and its dual, see [29, Theorem 4.1, Theorem 4.2]. We refer to [29] for details on the derivation of this facially reduced **SDP**.

We now provide the details for  $\tilde{V}$ , the gangster operator  $\mathcal{G}_{\hat{J}}$ , and the *gangster index set*,  $\hat{J}$ .

- Such choices of the matrix  $\tilde{V}$  are discussed in [19, 29].<sup>1</sup> To be specific, here we use the following  $V_j \in \mathbb{R}^{j \times (j-1)}$  that has full column rank with  $V_j^T e = 0$ ; set

$$V_j := \begin{bmatrix} I_{j-1} \\ -e_{j-1}^T \end{bmatrix}, \quad y := \frac{1}{n}(m \otimes e_n), \quad \tilde{V} := \begin{bmatrix} 1 & 0 \\ y & V_k \otimes V_n \end{bmatrix} \in \mathbb{R}^{(nk+1) \times ((k-1)(n-1)+1)}. \tag{1.7}$$

One can show that every feasible  $Y = \tilde{V} R \tilde{V}^T$  of the **SDP** relaxation (1.6) is contained in the following *minimal face*,  $\mathcal{F}$  of  $\mathbb{S}_+^{nk+1}$  with properties:

$$\mathcal{F} = \tilde{V} \mathbb{S}_+^{(n-1)(k-1)+1} \tilde{V}^T \trianglelefteq \mathbb{S}_+^{nk+1};$$

$$Y \in \mathcal{F} \implies \text{range}(Y) \subseteq \text{range}(\tilde{V}), \quad Y \in \text{relint}(\mathcal{F}) \implies \text{range}(Y) = \text{range}(\tilde{V}).$$

- We let  $\mathcal{G}_\Omega$  represent the *coordinate projection map* on  $\mathbb{S}^{nk+1}$  that chooses the elements in the index set  $\Omega$ , i.e.,

$$\mathcal{G}_\Omega(Y) = (Y_{ij})_{ij \in \Omega} \in \mathbb{R}^{|\Omega|}, \quad \Omega \subseteq \Delta_{0:nk} := \{ij : 0 \leq i \leq j \leq nk\}.$$

By abuse of notation, we assume that the (gangster)<sup>2</sup> indices are restricted to the upper triangular indices  $\Delta_{0:nk}$ , even when not specified so. We denote the complement of  $\Omega$  in  $\Delta_{0:nk}$  by  $\Omega^c$ . The adjoint of  $\mathcal{G}_\Omega$ , denoted by  $\mathcal{G}_\Omega^* : \mathbb{R}^{|\Omega|} \rightarrow \mathbb{S}^{nk+1}$ , is given by

$$(\mathcal{G}_\Omega^*(w))_{ij} = \begin{cases} \frac{1}{2} w_{ij} & \text{if } i \neq j \text{ and } ij \text{ or } ji \in \Omega, \\ w_{ii} & \text{if } i = j \text{ and } ij \in \Omega, \\ 0 & \text{otherwise.} \end{cases}$$

- The gangster index set  $\hat{J}$  is defined to be the union of the top left index (00) and the set of indices  $J$  corresponding to the diagonal elements in the off-diagonal blocks in  $\bar{Y}_{(ij)}$  for all  $i < j$  in (1.4).

<sup>1</sup> There are several ways of constructing such a matrix  $\tilde{V}$ . Another way is presented in (2.45), below.

<sup>2</sup> The name gangster refers to *shooting holes* in the matrix, a term coined originally by Philippe Toint.

## 2 SDP and DNN relaxations of MC

In this section, we show the details of how to derive a stronger **DNN** relaxation of **MC** with redundant constraints. First, in Sect. 2.1, we show that the gangster set  $\widehat{J}$  in the relaxation (1.6) can be enlarged when some of the set sizes  $m_i = 1$ . Then in Sect. 2.2 we derive a strengthened **SDP** relaxation (2.32) with the new larger gangster set. Next, in Sect. 2.3, we investigate some useful redundant constraints of the **SDP** relaxation (2.32); these constraints will be used for strengthening our algorithm in Sect. 3. Finally, in Sect. 2.4 we present our facially reduced **DNN** relaxation (2.48).

### 2.1 Gangster constraints

Proposition 2.1 shows an important observation about the gangster set.

**Proposition 2.1** *Let  $\mathcal{K} := \{1, \dots, k\}$ ,  $\mathcal{I} := \{i \in \mathcal{K} : m_i = 1\}$ , and the complement  $\mathcal{I}^c := \mathcal{K} \setminus \mathcal{I}$ . Define  $m_{\text{one}} \in \mathbb{R}^k$  by*

$$(m_{\text{one}})_i = \begin{cases} 1 & \text{if } i \in \mathcal{I}, \\ 0 & \text{if } i \in \mathcal{I}^c. \end{cases} \tag{2.1}$$

Let  $X \in \mathcal{M}_m$  and let  $x := \text{vec}(X)$ . Then the following hold:

$$[(E_k - I_k) \otimes I_n] \circ (xx^T) = 0 \quad \text{and} \quad [\text{Diag}(m_{\text{one}}) \otimes (E_n - I_n)] \circ (xx^T) = 0.$$

**Proof** The results can be easily obtained by manipulating with the constraints in set  $\mathcal{M}_m$ . □

Proposition 2.1 naturally gives rise to our definition of full gangster set  $J_0$ .

**Definition 2.2** (full gangster set,  $J_0$ ) Let  $m_{\text{one}}$  be defined in (2.1). We define  $J_0 \subseteq \Delta_{0:nk}$  to be the set of (gangster) indices corresponding to the ones in  $(E_k - I_k) \otimes I_n + \text{Diag}(m_{\text{one}}) \otimes (E_n - I_n)$ , i.e.,

$$J_0 := \Delta_{0:nk} \cap (\Theta_o \cup \Theta_{\mathcal{I}}), \tag{2.2}$$

where

$$\Theta_o := \{\text{all diagonal positions of all off-diagonal blocks in } \overline{Y}\},$$

$$\Theta_{\mathcal{I}} := \{\text{all off-diagonal positions of the } i\text{th diagonal blocks in } \overline{Y} \text{ if } m_i = 1\}.$$

Let  $\widehat{J}_0 := J_0 \cup (00)$ ,  $J_0$  in (2.2). Replacing the gangster set  $\widehat{J}$  in (1.6) by  $\widehat{J}_0$ , we obtain the following “strengthened” **SDP** relaxation immediately:

$$\begin{aligned}
 & \min \frac{1}{2} \operatorname{trace} \left( \tilde{V}^T L_A \tilde{V} \right) R \\
 \text{s.t. } & \mathcal{G}_{j_0}(\tilde{V}R\tilde{V}^T) = u_0 \\
 & R \geq 0.
 \end{aligned} \tag{2.3}$$

Note that when  $\mathcal{I} \neq \emptyset$ , the gangster set  $J_0$  is larger than  $J$  used in [19]. In particular, we see that if  $m_i = 1, \forall i \in \mathcal{K}$ , then necessarily all the diagonal elements of all off-diagonal blocks and all the off-diagonal elements of all diagonal blocks are zero. This is precisely the case for the quadratic assignment problem, **QAP**, e.g., [18, 31].

We indeed show below in Theorem 2.11 that the Slater condition, strict feasibility, holds for (2.3). It can also be shown that the singularity degree is one, i.e., only one step is needed to obtain strict feasibility.<sup>3</sup> However, there are redundant linear equality constraints in (2.3). Regarding this concern, the gangster constraint in (2.3) plays a crucial role. In this section, we study further properties of the gangster set  $J_0$  and the restricted gangster set  $J_{\mathcal{I}}$  defined in Definition 2.3. Then in Sect. 2.2, we present our **SDP** relaxation (2.32) (which uses  $J_{\mathcal{I}}$  in place of  $J_0$  in (2.3)) and establish some desirable regularity conditions. Specifically, we show that *Robinson regularity*<sup>4</sup> holds for (2.32) below.

Now, we carefully define the various parts of the restricted gangster set.

**Definition 2.3** (*Restricted gangster set,  $J_{\mathcal{I}}$* ) Let  $\mathcal{K} := \{1, \dots, k\}, \mathcal{I} := \{i \in \mathcal{K} : m_i = 1\}$ . Fix a  $j_0 \in \mathcal{I}^c$ .

Define the *gangster subsets*,  $J_i, i = 1, 2, 3$ , by

- $J_1 :=$  all diagonal positions of the  $(i, k)$  (and  $(k, i)$ ) blocks,  $\forall i \in \mathcal{I} \setminus \{k\}$ ;
- $J_2 :=$  all diagonal positions of the  $(j_0, k)$  (and  $(k, j_0)$ ) blocks;
- $J_3 :=$  all diagonal positions of the  $(k - 2, k - 1)$  (and  $(k - 1, k - 2)$ ) blocks.

Given  $J_0$  in (2.2), the *restricted gangster set*,  $J_{\mathcal{I}} \subseteq \Delta_{0:nk}$ , is:

$$J_{\mathcal{I}} = \begin{cases} J_0, & \text{if } \mathcal{I} = \emptyset \\ J_0 \setminus J_1, & \text{if } k \notin \mathcal{I} \neq \emptyset \\ J_0 \setminus (J_1 \cup J_2), & \text{if } k \in \mathcal{I} \neq \mathcal{K} \\ J_0 \setminus (J_1 \cup J_3), & \text{if } \mathcal{I} = \mathcal{K}. \end{cases} \tag{2.4}$$

<sup>3</sup> The singularity degree is essentially the minimum number of **FR** steps needed to find the so-called *minimal face*, the smallest face containing the feasible set. The singularity degree depends on the data of the problem, i.e., the linear constraints,  $\mathcal{A}(X) = b$ , and the semidefinite cone. For the original **SDP** relaxation before **FR**, it can be shown that the singularity degree is one, i.e., one can use the lifted linear equality constraints to find an exposing vector and use it to construct the matrix  $\hat{V}$ .

<sup>4</sup> Robinson regularity: strict feasibility holds and the linear constraints are onto, [23].



**Remark 2.4** Note that the restricted gangster set  $J_{\mathcal{I}}$  is obtained from  $J_0$  by removing some indices. We see below in Remark 2.9, that  $J_{\mathcal{I}}$  is in some sense the *largest effective subset* in  $J_0$ .

We next show in Theorem 2.6 below that the nullspaces of  $\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  and  $\mathcal{G}_{J_0}(\tilde{V} \cdot \tilde{V}^T)$  are the same. First, we recall the following.

**Lemma 2.5** ([29, Lemma 4.1]<sup>5</sup>) *Let  $R \in \mathbb{S}^{(n-1)(k-1)+1}$  be given,  $\tilde{V}$  be as in (1.7), and let*

$$Y = \tilde{V}R\tilde{V}^T.$$

Then the block notation of (1.4) yields:

$$m_i Y_{(j0)}^T = e^T \bar{Y}_{(ij)}, \quad \forall i, j \in \{1, \dots, k\}; \tag{2.5}$$

$$\sum_{i=1}^k \text{diag}(\bar{Y}_{(ij)}) = Y_{(j0)}, \quad \forall j \in \{1, \dots, k\}. \tag{2.6}$$

**Theorem 2.6** *Let  $J_0, J_{\mathcal{I}}, \tilde{V}$  be as in (2.2), (2.4), (1.7); and let  $Y = \tilde{V}R\tilde{V}^T$ , for some  $R \in \mathbb{S}^{(n-1)(k-1)+1}$ . Then*

$$\mathcal{G}_{J_{\mathcal{I}}}(Y) = 0 \iff \mathcal{G}_{J_0}(Y) = 0. \tag{2.7}$$

**Proof** The equivalence in (2.7) is trivially true if  $\mathcal{I} = \emptyset$ , since in this case necessarily  $J_{\mathcal{I}} = J_0$ . Thus, without loss of generality, we assume  $\mathcal{I} \neq \emptyset$ .

Since  $J_{\mathcal{I}} \subseteq J_0$ , we trivially have  $\mathcal{G}_{J_0}(Y) = 0 \implies \mathcal{G}_{J_{\mathcal{I}}}(Y) = 0$ . Hence, to establish (2.7), it remains to prove the converse implication, i.e., to show that

$$\mathcal{G}_{J_{\mathcal{I}}}(Y) = 0 \implies \mathcal{G}_{J_0}(Y) = 0. \tag{2.8}$$

In view of the definition of  $J_{\mathcal{I}}$ , to prove (2.8), it amounts to proving the following three implications:

$$\begin{cases} \mathcal{G}_{J_0 \setminus \omega_1}(Y) = 0 \implies \mathcal{G}_{J_1}(Y) = 0 & \text{if } k \notin \mathcal{I} \neq \emptyset; \\ \mathcal{G}_{J_0 \setminus (\omega_1 \cup \omega_2)}(Y) = 0 \implies \mathcal{G}_{J_1}(Y) = 0, \mathcal{G}_{J_2}(Y) = 0 & \text{if } k \in \mathcal{I} \neq \mathcal{K}; \\ \mathcal{G}_{J_0 \setminus (\omega_1 \cup \omega_3)}(Y) = 0 \implies \mathcal{G}_{J_1}(Y) = 0, \mathcal{G}_{J_3}(Y) = 0 & \text{if } \mathcal{I} = \mathcal{K}. \end{cases} \tag{2.9}$$

To prove these implications, we write  $Y$  in the block matrix form (1.4). Since  $m_i = 1, \forall i \in \mathcal{I}$ , from (2.5), we obtain  $Y_{(i0)}^T = e^T \bar{Y}_{(ii)}, \forall i \in \mathcal{I}$ . This, together with  $\mathcal{G}_{J_0 \setminus (\omega_1 \cup \omega_2 \cup \omega_3)}(Y) = 0$ , yields that

<sup>5</sup> There is a misprint error in [29, Lemma 4.1]: the variable  $Z$  in item (c) should be  $Y$ .

$$Y_{(i0)} = \text{diag}\left(\bar{Y}_{(ii)}\right), \quad \forall i \in \mathcal{I}. \quad (2.10)$$

We can now prove the first assertion in (2.9). Using (2.6) and  $\mathcal{G}_{J_0 \setminus J_1}(Y) = 0$ , we have

$$Y_{(j0)} = \text{diag}\left(\bar{Y}_{(jj)}\right) + \text{diag}\left(\bar{Y}_{(kj)}\right), \quad \forall j \in \mathcal{I} \setminus \{k\}.$$

Combining this with (2.10) and the symmetry of  $Y$ , we see that

$$\text{diag}\left(\bar{Y}_{(jk)}\right) = \text{diag}\left(\bar{Y}_{(kj)}\right) = 0, \quad \forall j \in \mathcal{I} \setminus \{k\}, \quad (2.11)$$

i.e.,  $\mathcal{G}_{J_1}(Y) = 0$ .

Next, we prove the second assertion in (2.9). The reasoning for  $\mathcal{G}_{J_1}(Y) = 0$  is the same as in the previous case. In addition, from  $\mathcal{G}_{J_0 \setminus (J_1 \cup J_2)}(Y) = 0$ , (2.11) and (2.6), we have

$$Y_{(k0)} = \text{diag}\left(\bar{Y}_{(j_0 k)}\right) + \text{diag}\left(\bar{Y}_{(kk)}\right).$$

Since  $k \in \mathcal{I}$ , from (2.10), we have

$$Y_{(k0)} = \text{diag}\left(\bar{Y}_{(kk)}\right).$$

In view of the above two equations and the symmetry of  $Y$ , we obtain

$$\text{diag}\left(\bar{Y}_{(kj_0)}\right) = \text{diag}\left(\bar{Y}_{(j_0 k)}\right) = 0,$$

i.e.,  $\mathcal{G}_{J_2}(Y) = 0$ .

Finally, we prove the third assertion in (2.9). It follows from (2.6) and  $\mathcal{G}_{J_0 \setminus (J_1 \cup J_3)}(Y) = 0$  that

$$Y_{(j0)} = \text{diag}\left(\bar{Y}_{(jj)}\right) + \text{diag}\left(\bar{Y}_{(kj)}\right), \quad \forall j \in \mathcal{I} \setminus \{k-2, k-1, k\}.$$

Together with (2.10) and the symmetry of  $Y$ , we have

$$\text{diag}\left(\bar{Y}_{(jk)}\right) = \text{diag}\left(\bar{Y}_{(kj)}\right) = 0, \quad \forall j \in \mathcal{I} \setminus \{k-2, k-1, k\}. \quad (2.12)$$

Combining this with (2.6) and  $\mathcal{G}_{J_0 \setminus (J_1 \cup J_3)}(Y) = 0$  gives

$$\begin{cases} \text{diag}\left(\bar{Y}_{(k-2 k-2)}\right) + \text{diag}\left(\bar{Y}_{(k-1 k-2)}\right) + \text{diag}\left(\bar{Y}_{(k k-2)}\right) = Y_{(k-2)} \\ \text{diag}\left(\bar{Y}_{(k-2 k-1)}\right) + \text{diag}\left(\bar{Y}_{(k-1 k-1)}\right) + \text{diag}\left(\bar{Y}_{(k k-1)}\right) = Y_{(k-1)} \\ \text{diag}\left(\bar{Y}_{(k-2 k)}\right) + \text{diag}\left(\bar{Y}_{(k-1 k)}\right) + \text{diag}\left(\bar{Y}_{(k k)}\right) = Y_{(k)} \end{cases}$$

Using this together with (2.10) and the symmetry of  $Y$ , we obtain

$$\begin{cases} \text{diag}(\bar{Y}_{(k-2k-1)}) + \text{diag}(\bar{Y}_{(k-2k)}) = 0 \\ \text{diag}(\bar{Y}_{(k-2k-1)}) + \text{diag}(\bar{Y}_{(k-1k)}) = 0 \\ \text{diag}(\bar{Y}_{(k-2k)}) + \text{diag}(\bar{Y}_{(k-1k)}) = 0 \end{cases}$$

Therefore, we have

$$\text{diag}(\bar{Y}_{(k-2k)}) = \text{diag}(\bar{Y}_{(k-1k)}) = \text{diag}(\bar{Y}_{(k-2k-1)}) = \text{diag}(\bar{Y}_{(k-1k-2)}) = 0.$$

This together with (2.12) yields that  $\mathcal{G}_{J_1}(Y) = 0$  and  $\mathcal{G}_{J_3}(Y) = 0$ . □

Next, in the following Theorem 2.8, we show that, together with the facial structure defined by  $\tilde{V} \cdot \tilde{V}^T$ , the gangster constraint only needs the smaller set  $J_{\mathcal{I}}$ , and the corresponding linear map is onto. First, we need the following lemma for use in Theorem 2.8.

**Lemma 2.7** *Let  $V_n$  be defined in (1.7), and let  $Z \in \mathbb{S}^n$ . If  $Z = \text{diag}(a)$  for some  $a \in \mathbb{R}^n$  or  $\text{diag}(Z) = 0$ , then we have*

$$V_n^T Z V_n = 0 \implies Z = 0.$$

**Proof**

Case 1: Let  $Z = \text{diag}(a) \in \mathbb{S}^n$ . Then

$$V_n^T Z V_n = \begin{bmatrix} a_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & a_{n-1} \end{bmatrix} + a_n E = 0 \implies a = 0 \implies Z = 0.$$

Case 2: Let  $Z \in \mathbb{S}^n$  with  $\text{diag}(Z) = 0$  be blocked as  $Z = \begin{bmatrix} C & b \\ b^T & 0 \end{bmatrix}$ , for some  $C \in \mathbb{S}^{n-1}$ ,  $\text{diag}(C) = 0$ , and  $b \in \mathbb{R}^{n-1}$ . Then

$$V_n^T Z V_n = C - eb^T - be^T = 0 \implies b = 0, C = 0 \implies Z = 0. \quad \square$$

We now show in Theorem 2.8 the onto property of the linear map defining the restricted gangster constraints,  $\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V}R\tilde{V}^T) = 0$ . A related result for the general graph partitioning problem but with another gangster set is given in [29, 30]. The basic idea is to show that the null space of its adjoint  $\tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(\cdot) \tilde{V}$  is zero.

**Theorem 2.8** *Let  $\tilde{V}$  be defined in (1.7) and  $J_{\mathcal{I}}$  be defined in (2.4). If  $w \in \mathbb{R}^{|\mathcal{I}|}$ , then*

$$\tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(w) \tilde{V} = 0 \implies w = 0.$$

**Proof** Let  $Y = \mathcal{G}_{J_{\mathcal{I}}}^*(w) \in \mathbb{S}^{nk+1}$ . Then we immediately have  $\tilde{V}^T Y \tilde{V} = 0$ . On the other hand, using the definition of  $\mathcal{G}_{J_{\mathcal{I}}}^*$ , we can block  $Y$  as  $Y = \begin{bmatrix} 0 & 0 \\ 0 & \bar{Y} \end{bmatrix}$ ,  $\bar{Y}$  as in (1.4), where  $\text{diag} \bar{Y}_{(ii)} = 0$ , and  $\bar{Y}_{(ij)}$  is diagonal whenever  $i \neq j$ . Let

$$Z := (V_k \otimes V_n)^T \bar{Y} (V_k \otimes V_n). \tag{2.13}$$

It follows from  $\tilde{V}^T Y \tilde{V} = 0$  that  $Z = 0$ . Note that

$$V_k \otimes V_n = \begin{bmatrix} V_n & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & V_n \\ -V_n & \cdots & -V_n \end{bmatrix}.$$

Therefore, if we write the above matrix  $Z$  in (2.13) as

$$\begin{bmatrix} Z_{(11)} & \cdots & Z_{(1\ k-1)} \\ \vdots & \ddots & \vdots \\ Z_{(k-1\ 1)} & \cdots & Z_{(k-1\ k-1)} \end{bmatrix},$$

we have

$$Z_{(ij)} = V_n^T \left( \bar{Y}_{(ij)} - \bar{Y}_{(kj)} - \bar{Y}_{(ik)} + \bar{Y}_{(kk)} \right) V_n = 0, \forall i, j \in \{1, \dots, k-1\}. \tag{2.14}$$

Furthermore, using the fact that  $\bar{Y}_{(ij)}$  is diagonal whenever  $i \neq j$ , we have

$$Z_{(ii)} = V_n^T \left( \bar{Y}_{(ii)} - 2\bar{Y}_{(ik)} + \bar{Y}_{(kk)} \right) V_n = 0, \forall i \in \{1, \dots, k-1\}. \tag{2.15}$$

It follows from (2.14) and (2.15) that

$$V_n^T \left( 2\bar{Y}_{(ij)} - \bar{Y}_{(ii)} - \bar{Y}_{(jj)} \right) V_n = 0, \forall i, j \in \{1, \dots, k-1\}. \tag{2.16}$$

We now claim that

$$\bar{Y}_{(ii)} = 0, \forall i \in \{1, \dots, k\}, \tag{2.17}$$

holds under the different choices of  $\mathcal{I}$  in  $J_{\mathcal{I}}$  given in (2.4).

- If  $\mathcal{I} = \emptyset$ , by (2.4), we have  $J_{\mathcal{I}} = J_0$ , i.e., (2.17) holds.
- If  $k \notin \mathcal{I} \neq \emptyset$ , then by (2.4), we have  $J_{\mathcal{I}} = J_0 \setminus J_1$ , i.e., the following equalities hold:

$$\bar{Y}_{(kk)} = 0 \tag{2.18}$$

$$\bar{Y}_{(ik)} = \bar{Y}_{(ki)} = 0, \forall i \in \mathcal{I} \tag{2.19}$$

$$\bar{Y}_{(ii)} = 0, \quad \forall i \in \{1, \dots, k-1\} \setminus \mathcal{I}. \tag{2.20}$$

From (2.15), (2.18), (2.19), we get  $V_n^T \bar{Y}_{(ii)} V_n = 0, \forall i \in \mathcal{I}$ . Notice that  $\bar{Y}_{(ii)}$  is a symmetric matrix with zeros on the diagonal, by Lemma 2.7, we get  $\bar{Y}_{(ii)} = 0, \forall i \in \mathcal{I}$ . This, together with (2.18) and (2.20), yields (2.17).

- If  $k \in \mathcal{I} \neq \mathcal{K}$ , then  $\mathcal{I}^c \neq \emptyset$ . By (2.4), we have  $J_{\mathcal{I}} = J_0 \setminus (J_1 \cup J_2)$ , i.e

$$\bar{Y}_{(ii)} = 0, \quad \forall i \in \mathcal{I}^c \tag{2.21}$$

$$\bar{Y}_{(kj_0)} = \bar{Y}_{(j_0k)} = 0, \quad \text{for the } j_0 \in \mathcal{I}^c \tag{2.22}$$

$$\bar{Y}_{(ki)} = \bar{Y}_{(ik)} = 0, \quad \forall i \in \mathcal{I} \setminus \{k\}. \tag{2.23}$$

It follows from (2.15), (2.21), (2.22), and Lemma 2.7 that

$$\bar{Y}_{(kk)} = 0. \tag{2.24}$$

In view of (2.15), (2.23), (2.24) and Lemma 2.7, we have  $\bar{Y}_{(ii)} = 0, \forall i \in \mathcal{I} \setminus \{k\}$ . This, together with (2.21) and (2.24), yields (2.17).

- If  $\mathcal{I} = \mathcal{K}$ , then by (2.4), we have  $J_{\mathcal{I}} = J_0 \setminus (J_1 \cup J_3)$ , i.e.,

$$\begin{aligned} \bar{Y}_{(k-1,k-2)} &= \bar{Y}_{(k-2,k-1)} = 0, \\ \bar{Y}_{(ki)} &= \bar{Y}_{(ik)} = 0, \forall i \in \{1, \dots, k-1\}. \end{aligned} \tag{2.25}$$

With  $i = k-1, j = k-2$  in (2.14), by (2.25) and Lemma 2.7, we have  $\bar{Y}_{(kk)} = 0$ . This together with (2.25), (2.15) and Lemma 2.7 yields (2.17).

In summary, the claim (2.17) holds. Combining (2.17) and (2.15), we get

$$V_n^T \bar{Y}_{(ki)} V_n = V_n^T \bar{Y}_{(ik)} V_n = 0 \quad \forall i \in \{1, \dots, k-1\}. \tag{2.26}$$

In addition, it follows from (2.17) and (2.16) that

$$V_n^T \bar{Y}_{(ij)} V_n = 0 \quad \forall i, j \in \{1, \dots, k-1\}. \tag{2.27}$$

Combining (2.26), (2.27) and (2.17), we have

$$V_n^T \bar{Y}_{(ij)} V_n = 0 \quad \forall i, j \in \{1, \dots, k\}.$$

Since  $\bar{Y}_{(ij)}$  is either a diagonal matrix or a matrix with diagonal equal to zeros, by Lemma 2.7 we have  $\bar{Y}_{(ij)} = 0$ , for all  $i, j \in \{1, \dots, k\}$ . Therefore,  $Y = 0$ . Thus, it follows that  $w = 0$ . □

**Remark 2.9** Combining Theorem 2.6 with Theorem 2.8, we see that the linear map  $\mathcal{G}_{J_0}(\tilde{V} \cdot \tilde{V}^T)$  is *not* onto but  $\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  is, and the two linear maps have the same nullspace. Since the restricted gangster set  $J_{\mathcal{I}}$  is obtained by removing indices in  $J_0$  and the linear map  $\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  is onto according to Theorem 2.8, this suggests that

we have removed *just the right number of indices* from  $J_0$ . By Theorem 2.6, all the indices in  $J_0 \setminus J_{\mathcal{I}}$  are redundant. Hence, we conclude that there does not exist a larger set that contains  $J_{\mathcal{I}}$  such that Theorem 2.8 holds, i.e., there are no redundant indices in  $J_{\mathcal{I}}$ .

We extend this result in Corollary 2.10 to show that the operator  $\mathcal{G}_{\hat{J}_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  is onto when considered as a linear transformation mapping into  $\mathbb{R}^{|\hat{J}_{\mathcal{I}}|+1}$ , where  $\hat{J}_{\mathcal{I}} := J_{\mathcal{I}} \cup \{00\}$  with  $J_{\mathcal{I}}$  defined in (2.4).

**Corollary 2.10** *Let  $\tilde{V}$  be as in (1.7),  $J_{\mathcal{I}}$  as in (2.4), and  $\hat{J}_{\mathcal{I}} := J_{\mathcal{I}} \cup \{00\}$ . If  $w \in \mathbb{R}^{|\hat{J}_{\mathcal{I}}|+1}$ , then*

$$\tilde{V}^T \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) \tilde{V} = 0 \implies w = 0.$$

Moreover, the linear transformation  $\mathcal{G}_{\hat{J}_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  is onto  $\mathbb{R}^{|\hat{J}_{\mathcal{I}}|+1}$ .

**Proof** For  $w \in \mathbb{R}^{|\hat{J}_{\mathcal{I}}|+1}$ , write  $w = [w_{00} \ \check{w}^T]^T$ , where  $\check{w} \in \mathbb{R}^{|\hat{J}_{\mathcal{I}}|}$ . Then we have

$$\mathcal{G}_{J_{\mathcal{I}}}^*(\check{w}) = \begin{bmatrix} 0 & 0 \\ 0 & \overline{W} \end{bmatrix} \quad \text{and} \quad \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) = \begin{bmatrix} w_{00} & 0 \\ 0 & \overline{W} \end{bmatrix}$$

for some  $\overline{W} \in \mathbb{S}^{n_k}$ . A direct computation using the definition of  $\tilde{V}$  yields

$$\tilde{V}^T \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) \tilde{V} = \begin{bmatrix} w_{00} + y^T \overline{W} y & y^T \overline{W} (V_k \otimes V_n) \\ (V_k^T \otimes V_n^T) \overline{W} y & (V_k^T \otimes V_n^T) \overline{W} (V_k \otimes V_n) \end{bmatrix}, \tag{2.28}$$

$$\tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(\check{w}) \tilde{V} = \begin{bmatrix} y^T \overline{W} y & y^T \overline{W} (V_k \otimes V_n) \\ (V_k^T \otimes V_n^T) \overline{W} y & (V_k^T \otimes V_n^T) \overline{W} (V_k \otimes V_n) \end{bmatrix}. \tag{2.29}$$

Now, assume that  $\tilde{V}^T \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) \tilde{V} = 0$ . Then we see from (2.28) that  $(V_k^T \otimes V_n^T) \overline{W} (V_k \otimes V_n) = 0$ . Following the same argument as in the proof of Theorem 2.8 (start from (2.13) and use  $\overline{W}$  in place of  $\overline{Y}$  there), we conclude that  $\overline{W} = 0$ . Combining this with (2.28) and the assumption  $\tilde{V}^T \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) \tilde{V} = 0$  gives

$$\begin{bmatrix} w_{00} & 0 \\ 0 & 0 \end{bmatrix} = \tilde{V}^T \mathcal{G}_{\hat{J}_{\mathcal{I}}}^*(w) \tilde{V} = 0,$$

showing that  $w_{00} = 0$ . On the other hand, we can deduce from (2.29) and the fact  $\overline{W} = 0$  that

$$\tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(\check{w}) \tilde{V} = 0.$$

This implies  $\check{w} = 0$ , according to Theorem 2.8. Consequently,  $w = [w_{00} \ \check{w}^T]^T = 0$ .

Note that when  $\mathcal{G}_{\tilde{J}_T}(\tilde{V} \cdot \tilde{V}^T)$  is considered as a linear transformation into  $\mathbb{R}^{|\mathcal{I}|+1}$ , its adjoint operator is  $\tilde{V}^T \mathcal{G}_{\tilde{J}_T}^*(\cdot) \tilde{V}$ . Since the kernel of  $\tilde{V}^T \mathcal{G}_{\tilde{J}_T}^*(\cdot) \tilde{V}$  is zero, we conclude that  $\mathcal{G}_{\tilde{J}_T}(\tilde{V} \cdot \tilde{V}^T)$  is onto. This completes the proof.  $\square$

### 2.2 Facially reduced SDP relaxation

Now we present our facially reduced **SDP** relaxation (2.32) (which uses  $\hat{J}_T$  in place of  $\hat{J}_0$  in (2.3)) and establish some desirable regularity conditions. Specifically, we show that *Robinson regularity*<sup>6</sup> holds for (2.32) below.

To obtain primal strict feasibility, we proceed as in [29, Theorem 4.1], and make use of the barycenter of the rank-1 matrices of the lifting [29, Equation (3.3), Theorem 3.1]):

$$\begin{aligned} \hat{Y} &:= \frac{m_1! \dots m_k!}{n!} \sum_{\text{Mat}(x) \in \mathcal{M}_n} \begin{bmatrix} 1 & x^T \\ x & xx^T \end{bmatrix} \\ &= \begin{bmatrix} 1 & \frac{m_1}{n} e_n^T & \dots & \frac{m_k}{n} e_n^T \\ \frac{m_1}{n} e_n & \left( \frac{m_1}{n} I_n + \frac{m_1(m_1-1)}{n(n-1)} (E_n - I_n) \right) & \dots & \left( \frac{m_1 m_k}{n(n-1)} \right) (E_n - I_n) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{m_k}{n} e_n & \left( \frac{m_1 m_k}{n(n-1)} \right) (E_n - I_n) & \dots & \left( \frac{m_k}{n} I_n + \frac{m_k(m_k-1)}{n(n-1)} (E_n - I_n) \right) \end{bmatrix}. \end{aligned} \tag{2.30}$$

On the other hand, to analyze dual strict feasibility, we define the following matrices:

$$\begin{aligned} \tilde{W} &:= \beta \begin{bmatrix} \alpha & 0 \\ 0 & 2Q_{\mathcal{I}} \end{bmatrix} \text{ and } Q_{\mathcal{I}} := T_{\mathcal{I}} \otimes I_n + S_{\mathcal{I}} \otimes (E_n - I_n), \quad \alpha < 0 < \beta; \\ (T_{\mathcal{I}}, S_{\mathcal{I}}) &= \begin{cases} (E_k - I_k, 0) & \text{if } \mathcal{I} = \emptyset, \\ (E_k - I_k - \hat{M}_{\text{one}}, e^T m_{\text{one}} M_{\text{one}}) & \text{if } k \notin \mathcal{I} \neq \emptyset, \\ (E_k - I_k - \hat{E}, M_{\text{one}}) & \text{if } k \in \mathcal{I} \neq \mathcal{K}, \\ (0, I_k) & \text{if } \mathcal{I} = \mathcal{K}, \end{cases} \end{aligned} \tag{2.31}$$

where  $m_{\text{one}}$ ,  $\mathcal{I}$  and  $\mathcal{K}$  are defined in Definition 2.3,

$$\hat{E} = \begin{bmatrix} 0 & e_{k-1} \\ e_{k-1}^T & 0 \end{bmatrix} \in \mathbb{S}^k, \quad M_{\text{one}} = \text{diag}(m_{\text{one}}), \quad \hat{M}_{\text{one}} = \begin{bmatrix} 0 & \hat{m}_{\text{one}} \\ \hat{m}_{\text{one}}^T & 0 \end{bmatrix} \in \mathbb{S}^k,$$

with  $\hat{m}_{\text{one}} \in \mathbb{R}^{k-1}$  being the vector that contains the first  $k - 1$  entries of  $m_{\text{one}}$ .

**Theorem 2.11** *The following holds:*

<sup>6</sup> Strict feasibility holds and the linear constraints are onto, [23].

1. The facially reduced SDP (2.3) is equivalent to the following equality constrained problem

$$\begin{aligned} \text{cut}(m) \geq p_{\text{SDP}}^* &= \min \frac{1}{2} \text{trace} \left( \tilde{V}^T L_A \tilde{V} \right) R \\ \text{s.t. } &\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V}R\tilde{V}^T) = \mathcal{G}_{J_{\mathcal{I}}}(u_0 u_0^T) \\ &R \succeq 0. \end{aligned} \tag{2.32}$$

2. The primal model (2.32) satisfies strict feasibility, with Slater point

$$\tilde{R} = \left[ \begin{array}{c|c} 1 & 0 \\ \hline 0 & \frac{1}{n^2(n-1)}(n \text{Diag}(\hat{m}_{k-1}) - \hat{m}_{k-1}\hat{m}_{k-1}^T) \otimes (nI_{n-1} - E_{n-1}) \end{array} \right] \in \mathbb{S}_{++}^{(k-1)(n-1)+1}, \tag{2.33}$$

where  $\hat{m}_{k-1} = (m_1, \dots, m_{k-1})^T \in \mathbb{Z}_+^{k-1}$ . Moreover,  $\tilde{V}\tilde{R}\tilde{V}^T = \hat{Y}$  given in (2.30); and Robinson regularity holds for (2.32).

3. The dual problem of (2.32) is

$$\begin{aligned} \max &\frac{1}{2}w_{00} \\ \text{s.t. } &\tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(w)\tilde{V} \leq \tilde{V}^T L_A \tilde{V}. \end{aligned} \tag{2.34}$$

Moreover, with  $\tilde{W}$  as in (2.31), and with sufficiently positive  $\beta$  and sufficiently negative  $\alpha$ , we get that the point  $\tilde{w}_{\mathcal{I}} := \mathcal{G}_{J_{\mathcal{I}}}(\tilde{W})$  is strictly feasible for (2.34).

**Proof** Item 1: By Theorem 2.6, the nullspaces of  $\mathcal{G}_{J_{\mathcal{I}}}(\tilde{V} \cdot \tilde{V}^T)$  and  $\mathcal{G}_{J_0}(\tilde{V} \cdot \tilde{V}^T)$  are the same. Therefore, the equivalence of (2.32) and (2.3) is obvious. Item 2: The strict feasibility of  $\tilde{R}$  follows immediately from [29, Theorem 4.1], which asserts  $\tilde{R} > 0$  and established  $\tilde{V}\tilde{R}\tilde{V}^T = \hat{Y}$  in its proof. The Robinson regularity holds in view of the strict feasibility of  $\tilde{R}$  and Theorem 2.8.

Item 3: It is standard to show that the dual problem of (2.32) is given by (2.34). We now prove the claim concerning strict feasibility.

With the  $y$  in (1.7), the  $\tilde{V}$  in (1.7), the definitions of  $\tilde{W}$  and  $\tilde{w}_{\mathcal{I}}$ , and the definition of  $J_{\mathcal{I}}$  in Definition 2.3, we can compute that

$$\begin{aligned} \tilde{V}^T \mathcal{G}_{J_{\mathcal{I}}}^*(\tilde{w}_{\mathcal{I}})\tilde{V} &= \beta \begin{bmatrix} 1 & y^T \\ 0 & V_k^T \otimes V_n^T \end{bmatrix} \begin{bmatrix} \alpha & 0 \\ 0 & Q_{\mathcal{I}} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ y & V_k \otimes V_n \end{bmatrix} \\ &= \beta \begin{bmatrix} \alpha + y^T Q_{\mathcal{I}} y & y^T Q_{\mathcal{I}} (V_k \otimes V_n) \\ (V_k^T \otimes V_n^T) Q_{\mathcal{I}} y & (V_k^T \otimes V_n^T) Q_{\mathcal{I}} (V_k \otimes V_n) \end{bmatrix}. \end{aligned} \tag{2.35}$$

Now, recall the following relations, which are immediate consequences of the definition of  $V_j$ :

$$V_j^T = [I_{j-1} \quad -e_{j-1}], \quad V_j^T E_j = V_j^T e_j e_j^T = 0, \quad \text{and} \quad V_j^T V_j = E_{j-1} + I_{j-1}.$$

Then we have



$$\begin{aligned}
 (V_k^T \otimes V_n^T)Q_{\mathcal{I}}y &= (V_k^T \otimes V_n^T)(T_{\mathcal{I}} \otimes I_n + S_{\mathcal{I}} \otimes (E_n - I_n))y \\
 &= (V_k^T T_{\mathcal{I}} \otimes V_n^T + V_k^T S_{\mathcal{I}} \otimes V_n^T (E_n - I_n))y \\
 &= (V_k^T T_{\mathcal{I}} \otimes V_n^T - V_k^T S_{\mathcal{I}} \otimes V_n^T)y \\
 &= (V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}}) \otimes V_n^T)(m \otimes e_n)/n \\
 &= (V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}})m) \otimes V_n^T e_n/n = 0
 \end{aligned}$$

and

$$\begin{aligned}
 (V_k^T \otimes V_n^T)Q_{\mathcal{I}}(V_k \otimes V_n) &= (V_k^T \otimes V_n^T)(T_{\mathcal{I}} \otimes I_n + S_{\mathcal{I}} \otimes (E_n - I_n))(V_k \otimes V_n) \\
 &= V_k^T T_{\mathcal{I}} V_k \otimes V_n^T V_n + V_k^T S_{\mathcal{I}} V_k \otimes V_n^T (E_n - I_n) V_n \\
 &= V_k^T T_{\mathcal{I}} V_k \otimes V_n^T V_n - V_k^T S_{\mathcal{I}} V_k \otimes V_n^T V_n \\
 &= V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}}) V_k \otimes V_n^T V_n \\
 &= V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}}) V_k \otimes (I_{n-1} + E_{n-1}).
 \end{aligned}$$

Combining the above two displays with (2.35), we obtain

$$\tilde{V}^T \mathcal{G}_{\mathcal{I}}^* (\tilde{w}_{\mathcal{I}}) \tilde{V} = \beta \begin{bmatrix} \alpha + y^T Q_{\mathcal{I}} y & 0 \\ 0 & V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}}) V_k \otimes (I_{n-1} + E_{n-1}) \end{bmatrix}. \tag{2.36}$$

We next show that  $V_k(T_{\mathcal{I}} - S_{\mathcal{I}})V_k < 0$  in each of the four cases in the definition of  $J_{\mathcal{I}}$ .

- If  $\mathcal{I} = \emptyset$ , then  $V_k^T(T_{\mathcal{I}} - S_{\mathcal{I}})V_k = V_k^T(E_k - I_k)V_k = -V_k^T V_k = -(I_{k-1} + E_{k-1}) < 0$ ,
- If  $k \notin \mathcal{I} \neq \emptyset$ , then we have

$$\begin{aligned}
 V_k^T (T_{\mathcal{I}} - S_{\mathcal{I}}) V_k &= V_k^T (E_k - I_k - \hat{M}_{\text{one}} - e^T m_{\text{one}} M_{\text{one}}) V_k \\
 &= -I_{k-1} - E_{k-1} - V_k^T (\hat{M}_{\text{one}} + e^T m_{\text{one}} M_{\text{one}}) V_k \\
 &\leq -I_{k-1} - E_{k-1} - V_k^T (\hat{M}_{\text{one}} + m_{\text{one}} m_{\text{one}}^T) V_k \\
 &= -I_{k-1} - E_{k-1} - V_k^T \left( \begin{bmatrix} 0 & \hat{m}_{\text{one}} \\ \hat{m}_{\text{one}}^T & 0 \end{bmatrix} + \begin{bmatrix} \hat{m}_{\text{one}} \\ 0 \end{bmatrix} \begin{bmatrix} \hat{m}_{\text{one}}^T & 0 \end{bmatrix} \right) V_k \\
 &= -I_{k-1} - E_{k-1} - [I_{k-1} \quad -e_{k-1}] \begin{bmatrix} \hat{m}_{\text{one}} \hat{m}_{\text{one}}^T & \hat{m}_{\text{one}} \\ \hat{m}_{\text{one}}^T & 0 \end{bmatrix} \begin{bmatrix} I_{k-1} \\ -e_{k-1}^T \end{bmatrix} \\
 &= -I_{k-1} - E_{k-1} - (\hat{m}_{\text{one}} \hat{m}_{\text{one}}^T - e_{k-1} \hat{m}_{\text{one}}^T - \hat{m}_{\text{one}} e_{k-1}^T) \\
 &= -I_{k-1} - (e_{k-1} - \hat{m}_{\text{one}})(e_{k-1} - \hat{m}_{\text{one}})^T \\
 &\leq -I_{k-1} < 0,
 \end{aligned}$$

where the first “ $\leq$ ” follows from the observation that  $e^T m_{\text{one}} M_{\text{one}} \geq m_{\text{one}} m_{\text{one}}^T$ .

- If  $k \in \mathcal{I} \neq \emptyset$ , then we have

$$\begin{aligned}
 V_k^T(T_{\mathcal{I}} - S_{\mathcal{I}})V_k &= V_k^T(E_k - I_k - \widehat{E} - M_{\text{one}})V_k \\
 &= -I_{k-1} - E_{k-1} - V_k^T(\widehat{E} + M_{\text{one}})V_k \\
 &= -I_{k-1} - E_{k-1} - \begin{bmatrix} I_{k-1} & -e_{k-1} \end{bmatrix} \begin{bmatrix} \text{diag}(\widehat{m}_{\text{one}}) & e \\ e^T & 1 \end{bmatrix} \begin{bmatrix} I_{k-1} \\ -e^T \end{bmatrix} \\
 &= -I_{k-1} - E_{k-1} - (\text{diag}(\widehat{m}_{\text{one}}) - E_{k-1}) \\
 &= -I_{k-1} - \text{diag}(\widehat{m}_{\text{one}}) < 0
 \end{aligned}$$

- If  $\mathcal{I} = \mathcal{K}$ , then we have  $V_k^T(T_{\mathcal{I}} - S_{\mathcal{I}})V_k = V_k^T(-I_k)V_k = -(E_{k-1} + I_{k-1}) < 0$ .

In summary, we have  $V_k(T_{\mathcal{I}} - S_{\mathcal{I}})V_k < 0$ , which together with  $I_{n-1} + E_{n-1} > 0$  yields that  $V_k^T(T_{\mathcal{I}} - S_{\mathcal{I}})V_k \otimes (I_{n-1} + E_{n-1}) < 0$  in (2.36). Therefore, with  $\alpha \ll 0 \ll \beta$ , we have  $\widehat{V}^T \mathcal{G}_{\mathcal{I}}^*(\widetilde{w}_{\mathcal{I}})\widehat{V} \leq \widehat{V}^T L_A \widehat{V}$ , i.e.,  $\widetilde{w}_{\mathcal{I}}$  is strictly feasible for (2.34).  $\square$

We emphasize that (2.32) is a **SDP** relaxation of model (1.2). It uses facial reduction to guarantee strict feasibility for both the relaxation and its dual. The Robinson regularity condition holds and thus we obtain robustness.

### 2.3 Redundant constraints of the SDP relaxation

In this section, we identify important redundant constraints of (2.32). These redundant constraints are used in the numerical implementations in Sect. 3, below. We first define some useful linear transformations on  $\mathbb{S}^{nk+1}$ .

**Definition 2.12** Let  $Y \in \mathbb{S}^{nk+1}$  be blocked as in (1.4). Define the linear transformations on  $\mathbb{S}^{nk+1}$ :

1. arrow :

$$\text{arrow}(Y) := \text{diag}(Y) - \begin{bmatrix} 0 \\ Y_{1:nk0} \end{bmatrix} \in \mathbb{R}^{nk+1}.$$

2.  $\mathcal{D}_t, \mathcal{D}_d$ , and  $\mathcal{D}_o$  :

$$\begin{aligned}
 \mathcal{D}_t(Y) &:= \left( \text{trace } \overline{Y}_{(ij)} \right) \in \mathbb{S}^k; & \mathcal{D}_d(Y) &:= \sum_{i=1}^k \text{diag } \overline{Y}_{(ii)} \in \mathbb{R}^n; \\
 \mathcal{D}_o(Y) &:= \left( \sum_{s \neq t} \left( \overline{Y}_{(ij)} \right)_{st} \right) \in \mathbb{S}^k.
 \end{aligned}$$

The following Theorem 2.13 shows that the constraints in (2.37) and (2.38) below based on the linear transformations arrow,  $\mathcal{D}_t$ ,  $\mathcal{D}_d$ , and  $\mathcal{D}_o$  are redundant for the **SDP** relaxation (2.32). Though redundant, these constraints are useful for strengthening the subproblems of the **sPRSM** in Sect. 3.

**Theorem 2.13** Let  $Y = \widetilde{V}R\widetilde{V}^T$ , where  $R$  is feasible for (2.32), and  $\widehat{M} := mm^T - M$ . Then the following holds:

$$\text{arrow}(Y) = u_0; \tag{2.37}$$

$$\mathcal{D}_t(Y) = M; \quad \mathcal{D}_d(Y) = e_n; \quad \mathcal{D}_o(Y) = \widehat{M}. \tag{2.38}$$

**Proof** Note that in (2.32), it holds that  $\mathcal{G}_{J_T}(Y) = 0$ . Thus by Theorem 2.6, we have  $\mathcal{G}_{J_0}(Y) = 0$ , i.e., all the diagonal elements of off-diagonal blocks of  $\bar{Y}$  (see the block structure in (2.38)) are zero. From (2.6) in Lemma 2.5, we have that (2.37) holds. Also, as shown in [19, Theorem 5.1], the gangster constraint  $\mathcal{G}_{J_0}(Y) = 0$  together with  $Y_{00} = 1$  and  $R \geq 0$  shows that  $Y = \widetilde{V}R\widetilde{V}^T$  satisfies all the constraints in (2.38) except for

$$\mathcal{D}_o(Y) = \widehat{M}. \tag{2.39}$$

Now it suffices to show that  $Y$  satisfies (2.39). Let  $D_2$  be defined as

$$D_2 := \begin{bmatrix} m^T m & -m^T \otimes e_n^T \\ -m \otimes e_n & I_k \otimes (e_n e_n^T) \end{bmatrix} = \begin{bmatrix} -m^T \\ I_k \otimes e_n \end{bmatrix} \begin{bmatrix} -m^T \\ I_k \otimes e_n \end{bmatrix}^T \geq 0.$$

Recall that  $\widetilde{V}$  is selected such that  $\text{range}(\widetilde{V}) \subseteq \text{null}(D_2)$  in [19, Page 351]. Thus  $\text{trace}(D_2 Y) = 0$ . Since  $R \geq 0$  and  $\mathcal{G}_{J_T}(\widetilde{V}R\widetilde{V}^T) = u_0$ , we have  $Y \geq 0$ ,  $Y_{00} = 1$ . Let  $v_1 = [1 \ Y_{1:nk0}^T]^T$ . Then we have

$$\begin{aligned} Y - v_1 v_1^T &= \begin{bmatrix} 1 & Y_{1:nk0}^T \\ Y_{1:nk0} & \bar{Y} \end{bmatrix} - \begin{bmatrix} 1 \\ Y_{1:nk0} \end{bmatrix} \begin{bmatrix} 1 & Y_{1:nk0} \end{bmatrix}^T \\ &= \begin{bmatrix} 0 & 0 \\ 0 & \bar{Y} - Y_{1:nk0} Y_{1:nk0}^T \end{bmatrix}. \end{aligned} \tag{2.40}$$

Note that  $\bar{Y} - Y_{1:nk0} Y_{1:nk0}^T$  is the Schur complement of  $Y_{00}$  in  $Y$  and  $Y \geq 0$ . Hence, it holds that  $\bar{Y} - Y_{1:nk0} Y_{1:nk0}^T \geq 0$ . Consequently, we deduce from (2.40) that  $Y \geq v_1 v_1^T$ .

Let  $X = \text{Mat}(Y_{1:nk0})$ . Since

$$\text{trace } D_2 Y = 0, \quad D_2 \geq 0, \quad \text{and } Y \geq v_1 v_1^T,$$

we see that

$$0 = \text{trace}(D_2 Y) \geq \text{trace}(D_2 v_1 v_1^T) = \|X^T e - m\|^2 \quad \text{and} \quad Y \begin{bmatrix} -m^T \\ I_k \otimes e_n \end{bmatrix} = 0. \tag{2.41}$$

Using the second relation in (2.41) together with the block partition of  $Y$  in (1.4), we have

$$-Y_{1:nk0} m^T + \bar{Y}(I_k \otimes e_n) = 0.$$

Multiplying the above relation on the left by  $I_k \otimes e_n^T$ , we obtain further that

$$-(I_k \otimes e_n^T)Y_{1:nk0}m^T + (I_k \otimes e_n^T)\bar{Y}(I_k \otimes e_n) = 0. \tag{2.42}$$

Next, recall from the first relation in (2.41) that  $(I_k \otimes e_n^T)Y_{1:nk0} = X^T e_n = m$ . Moreover, a direct computation shows that  $(I_k \otimes e_n^T)\bar{Y}(I_k \otimes e_n) = (e_n^T \bar{Y}_{(ij)} e_n)$ . Combining these with (2.42) yields

$$(e_n^T \bar{Y}_{(ij)} e_n) = mm^T.$$

Finally, it follows from the above equation and  $\mathcal{D}_t(Y) = M$  in (2.38) that

$$\mathcal{D}_o(Y) = \left( \sum_{s \neq t} (\bar{Y}_{(ij)})_{st} \right) = (e_n^T \bar{Y}_{(ij)} e_n) - \mathcal{D}_t(Y) = mm^T - M = \hat{M}. \quad \square$$

In Corollary 2.14 we provide a useful shift for our implementation of the objective function of (2.32).

**Corollary 2.14** *Let  $Y = \tilde{V}R\tilde{V}^T$ , where  $R$  is feasible for (2.32). Partition  $Y$  in blocks as in (1.4). Then we have*

$$\text{trace } Y = n + 1, \quad e^T Y_{(i0)} = m_i, \quad i = 1, \dots, k. \tag{2.43}$$

Moreover, the objective value in (1.5) satisfies

$$\text{trace}(L_A + \alpha I)Y = \text{trace } L_A Y + \alpha(n + 1), \quad \forall \alpha \in \mathbb{R}. \tag{2.44}$$

**Proof** It follows from (2.37),  $\mathcal{D}_t(Y) = M$  in (2.38) and  $m^T e = n$  that both constraints in (2.43) hold. Thus, the equality (2.44) holds evidently.  $\square$

### 2.4 DNN relaxation

For our **DNN** relaxation and algorithm in Sect. 3, below, we need the following matrix  $\hat{V}$  that have orthonormal columns.

**Assumption 2.15** Without loss of generality, by using a QR or SVD factorization on  $\tilde{V}$  in (1.7), or some other special construction if needed, we assume that the columns of  $\hat{V}$  form an orthonormal basis for the range of  $\tilde{V}$ . One such choice of  $\hat{V}$  is

$$\hat{V} = \begin{bmatrix} s & 0 \\ s_y & \hat{V}_k \otimes \hat{V}_n \end{bmatrix}, \tag{2.45}$$

where  $s := \sqrt{\frac{n}{n + \|m\|^2}}$  with  $\|m\|$  denoting the  $\ell_2$  norm of  $m$ , and  $\hat{V}_j$  is a matrix with orthonormal columns that satisfies  $\hat{V}_j^T e_j = 0$ .

Since the range of  $\widehat{V}$  is the same as the range of  $\widetilde{V}$ , we obtain the same minimal face

$$\widehat{V}\mathbb{S}_+^{(k-1)(n-1)+1}\widehat{V}^T = \widetilde{V}\mathbb{S}_+^{(k-1)(n-1)+1}\widetilde{V}^T.$$

Using  $\widehat{V}$  in place of  $\widetilde{V}$ , the facially reduced **SDP** (2.32) can be equivalently written as

$$\begin{aligned} \text{cut}(m) \geq p_{\text{SDP}}^* &= \min \frac{1}{2} \text{trace} \left( \widehat{V}^T L_A \widehat{V} \right) R \\ \text{s.t. } &\mathcal{G}_{\widehat{J}_T}(\widehat{V}R\widehat{V}^T) = \mathcal{G}_{\widehat{J}_T}(u_0 u_0^T) \\ &R \geq 0. \end{aligned} \tag{2.46}$$

The dual problem of (2.46) is

$$\begin{aligned} \max &\frac{1}{2} w_{00} \\ \text{s.t. } &\widehat{V}^T \mathcal{G}_{\widehat{J}_T}^*(w) \widehat{V} \leq \widehat{V}^T L_A \widehat{V}. \end{aligned} \tag{2.47}$$

The **SDP** relaxation (2.46) can be further strengthened by adding additional constraints. With the additional nonnegativity box constraint  $0 \leq (\widehat{V}R\widehat{V}^T)_{\widehat{J}_0^c} \leq 1$ , where  $\widehat{J}_0^c$  is the complement of  $\widehat{J}_0$ , we obtain the following doubly nonnegative, **DNN**, relaxation,

$$\begin{aligned} \text{cut}(m) \geq p_{\text{DNN}}^* &= \min \frac{1}{2} \text{trace} \left( \widehat{V}^T L_A \widehat{V} \right) R \\ \text{s.t. } &\mathcal{G}_{\widehat{J}_T}(\widehat{V}R\widehat{V}^T) = \mathcal{G}_{\widehat{J}_T}(u_0 u_0^T) \\ &R \geq 0 \\ &0 \leq \left( \widehat{V}R\widehat{V}^T \right)_{\widehat{J}_0^c} \leq 1. \end{aligned} \tag{2.48}$$

Note that the term **DNN** refers to the two cones constraints in (2.48), i.e., the positive semidefinite cone and the nonnegative cone. By Theorem 2.8 and Theorem 2.6, there are no redundant constraints in (2.48). Therefore, model (2.48) is somewhat the “most simplified” **DNN** relaxation of the **MC** problem.

The following Proposition 2.16 shows that the Slater point  $\widetilde{w}_T$  for (2.34) found in Theorem 2.11 is still strictly feasible for (2.47). Moreover, starting from the Slater point  $\widetilde{R}$  in (2.33) for (2.32), one can construct a Slater point for both (2.46) and (2.48): the fact that (2.48) has a Slater point will be important for our algorithmic development later.

**Theorem 2.16** *The strictly feasible point  $\widetilde{w}_T$  for (2.34) found in Theorem 2.11 is strictly feasible for (2.47). Moreover, define*

$$\widehat{R} := \widehat{V}^\dagger \widetilde{V} \widetilde{R} \widetilde{V}^T (\widehat{V}^\dagger)^T, \tag{2.49}$$

where  $\widetilde{R}$  is defined in (2.33),  $\widehat{V}^\dagger$  is the pseudoinverse of  $\widehat{V}$ , and  $\widetilde{V}$  and  $\widehat{V}$  are given in (1.7) and (2.45), respectively. Then it holds that  $\widehat{R}$  is strictly feasible for both (2.46) and (2.48), and  $\widehat{V} \widehat{R} \widehat{V}^T = \widehat{Y}$ , where  $\widehat{Y}$  is defined in (2.30).

**Proof**

1. Note that  $\text{Range}(\widehat{V}) = \text{Range}(\widetilde{V})$  by construction. This implies that  $\widehat{V}\widehat{V}^\dagger\widetilde{V} = \widetilde{V}$ . Thus, we have

$$\widetilde{V}^T(\widehat{V}^T)^\dagger\widehat{V}^T(L_A - \mathcal{G}_{J_T}^*(\widetilde{w}_T))\widehat{V}\widehat{V}^\dagger\widetilde{V} = \widetilde{V}^T(L_A - \mathcal{G}_{J_T}^*(\widetilde{w}_T))\widetilde{V} > 0,$$

where the positive definiteness follows from the fact that  $\widetilde{w}_T$  is strictly feasible for (2.34). Since  $(\widehat{V}^\dagger\widehat{V})^T = \widetilde{V}^T(\widehat{V}^T)^\dagger$  is a square matrix, we conclude from the above display that the matrix  $\widetilde{V}^T(\widehat{V}^T)^\dagger$  is nonsingular. Thus, we deduce further that

$$\widehat{V}^T(L_A - \mathcal{G}_{J_T}^*(\widetilde{w}_T))\widehat{V} = [\widetilde{V}^T(\widehat{V}^T)^\dagger]^{-1}\widetilde{V}^T(L_A - \mathcal{G}_{J_T}^*(\widetilde{w}_T))\widetilde{V}[\widehat{V}^\dagger\widetilde{V}]^{-1} > 0,$$

i.e.,  $\widetilde{w}_T$  is strictly feasible for (2.47).

2. The positive definiteness of  $\widehat{R}$  follows immediately from the fact that  $\widetilde{R} > 0$  (see Theorem 2.11 Item 2:) and the nonsingularity of  $\widetilde{V}^T(\widehat{V}^T)^\dagger$  just established. In addition, since  $\text{Range}(\widehat{V}) = \text{Range}(\widetilde{V})$ , we have  $\widehat{V}\widehat{V}^\dagger\widetilde{V} = \widetilde{V}$ . Using this and the definition of  $\widehat{R}$ , we see further that

$$\widehat{V}\widehat{R}\widehat{V}^T = \widehat{V}\widehat{V}^\dagger\widetilde{V}\widetilde{R}\widetilde{V}^T(\widehat{V}^\dagger)^T\widehat{V}^T = \widetilde{V}\widetilde{R}\widetilde{V}^T = \widehat{Y},$$

where the last equality follows from Theorem 2.11 Item 2. Then we obtain immediately that  $\mathcal{G}_{J_T}(\widehat{V}\widehat{R}\widehat{V}^T) = \mathcal{G}_{J_T}(\widehat{Y}) = 0$ . Consequently,  $\widehat{R}$  is strictly feasible for (2.46).

Finally, notice that entries of  $\widehat{Y}$  in  $J_0^c$  are strictly positive and strictly less than 1. Hence, we also have  $0 < (\widehat{V}\widehat{R}\widehat{V}^T)_{J_0^c} < 1$ . Thus, we have shown that  $\widehat{R}$  is strictly feasible for (2.48) and  $\widehat{V}\widehat{R}\widehat{V}^T = \widehat{Y}$ . □

The **DNN** problem (2.48) is extremely difficult for interior point methods, especially when the dimension is large. Motivated by the recent success in the application of splitting methods to the quadratic assignment problem in [18], we adopt a similar approach here. We first introduce a new variable and add the constraint  $Y = \widehat{V}\widehat{R}\widehat{V}^T$  to (2.48). By doing so, we essentially double the number of variables and transform the original problem (2.48) to the following equivalent model,

$$\begin{aligned} p_{\text{DNN}}^* &= \min \frac{1}{2} \text{trace } L_A Y \\ \text{s.t. } & Y = \widehat{V}\widehat{R}\widehat{V}^T \\ & \mathcal{G}_{J_T}(Y) = \mathcal{G}_{J_T}(u_0 u_0^T) \\ & R \geq 0 \\ & 0 \leq \mathcal{G}_{J_0^c}(Y) \leq 1. \end{aligned} \tag{2.50}$$

This is a separable convex programming problem with linear coupling constraints from the facial reduction. One can then apply first order splitting methods, which

allows us to take advantage of the two variables and the two cones to obtain two separate subproblems. We discuss one such method and the efficient solutions of the corresponding subproblems in closed form, in Sect. 3 below.

In passing, we would like to emphasize that the problem (2.50) is stable in that it has no redundant equality constraints, even though we added an extra linear constraint and a new variable  $Y$ . In detail, let  $\mathcal{T} : \mathbb{S}^{nk+1} \times \mathbb{S}^{(n-1)(k-1)+1} \rightarrow \mathbb{S}^{nk+1} \times \mathbb{R}^{|J_I|+1}$  be the linear operator defined as

$$\mathcal{T}(Y, R) = \begin{bmatrix} Y - \widehat{V}R\widehat{V}^T \\ \mathcal{G}_{J_I}^*(Y) \end{bmatrix}, \tag{2.51}$$

where  $\widehat{V}$  is defined in (2.45). Then its adjoint operator is

$$\mathcal{T}^*(W, w) = \begin{bmatrix} W + \mathcal{G}_{J_I}^*(w) \\ -\widehat{V}^T W \widehat{V} \end{bmatrix}.$$

We show in Proposition 2.17 below, that the operator  $\mathcal{T}$  is an *onto* linear transformation.

**Proposition 2.17**

1. Suppose that  $\mathcal{T}$  is given in (2.51) and  $(W, w) \in \mathbb{S}^{nk+1} \times \mathbb{R}^{|J_I|+1}$ . Then

$$\mathcal{T}^*(W, w) = 0 \implies (W, w) = 0.$$

2. Primal Slater points of model (2.50) are given by  $\widehat{R}$  in (2.49) and  $\widehat{Y}$  in (2.30).

**Proof**

1. Algebraic manipulation of  $\mathcal{T}^*(W, w) = 0$  yields the following two equations,

$$W + \mathcal{G}_{J_I}^*(w) = 0 \quad \text{and} \quad \widehat{V}^T W \widehat{V} = 0. \tag{2.52}$$

Combining the above two equations, we have  $\widehat{V}^T \mathcal{G}_{J_I}^*(w) \widehat{V} = 0$ . This implies that

$$\widetilde{V}^T (\widehat{V}^T)^\dagger \widehat{V}^T \mathcal{G}_{J_I}^*(w) \widehat{V} \widehat{V}^\dagger \widetilde{V} = 0.$$

Next, recall that  $\text{Range}(\widehat{V}) = \text{Range}(\widetilde{V})$  by construction. Thus, we have  $\widehat{V} \widehat{V}^\dagger \widetilde{V} = \widetilde{V}$ . Combining this with the above display yields  $\widetilde{V}^T \mathcal{G}_{J_I}^*(w) \widetilde{V} = 0$ . Then we deduce from Theorem 2.8 that  $w = 0$ . This together with the first relation in (2.52) gives  $W = 0$  and completes the proof.

2. This follows immediately from Proposition 2.16. □

### 3 sPRSM for DNN relaxation

In this section, we adapt the P-R splitting method [11] to solving our **DNN** relaxation (2.50). In essence, we separate the semidefinite cone constraints from the polyhedral constraints and obtain two subproblems. However, we also add back some provably redundant constraints. This is because these constraints are *not* redundant when the subproblems are considered as *independent optimization problems*. We take advantage of this and bring a constraint back if it does not increase the computational cost excessively. We denote this new method by **FRSMR**.

#### 3.1 FRSMR, A facially reduced splitting method with redundancies

Let  $L_s := \frac{1}{2}L_A$ . We can clearly rewrite (2.50) as

$$p_{\text{DNN}}^* = \min \text{ trace } L_s Y + \mathbb{1}_{\mathcal{Y}_o}(Y) + \mathbb{1}_{\mathcal{R}_o}(R) \tag{3.1}$$

$$\text{s.t. } Y = \widehat{V}R\widehat{V}^T.$$

where we use the *indicator function*,  $\mathbb{1}_{\mathcal{S}}(\mathcal{S})$ , that takes the value 0 on the set  $\mathcal{S}$  and  $\infty$  outside of  $\mathcal{S}$ ; and the two constraint sets in (3.1) are

$$\mathcal{R}_o := \mathbb{S}_+^{(k-1)(n-1)+1}, \quad \mathcal{Y}_o := \{Y \in \mathbb{S}^{nk+1} : \mathcal{G}_{\widehat{J}_X}(Y) = \mathcal{G}_{\widehat{J}_X}(u_0 u_0^T), 0 \leq \mathcal{G}_{\widehat{J}_0}(Y) \leq 1\}. \tag{3.2}$$

While this trivial decomposition is intuitive, a *splitting* method might benefit by operating on *tighter* constraint sets in the variables  $R$  and  $Y$ . Here, we shrink the sets in (3.2) by adding the following redundant constraints to (3.1):

1.  $\text{trace } R = n + 1$ , whose redundancy follows from Corollary 2.14,  $Y = \widehat{V}R\widehat{V}^T$  and  $\widehat{V}^T\widehat{V} = I$ .
2.  $\mathcal{D}_o(Y) = \widehat{M}$ , whose redundancy follows from Theorem 2.13.
3.  $\mathcal{G}_{\widehat{J}_0 \setminus \widehat{J}_X}(Y) = \mathcal{G}_{\widehat{J}_0 \setminus \widehat{J}_X}(u_0 u_0^T)$ , whose redundancy follows from Theorem 2.6.
4.  $e^T Y_{(i0)} = m_i$  for  $i = 1, \dots, k$ , whose redundancy follows from Corollary 2.14.

We thus arrive at the following equivalent problem of (3.1):

$$p_{\text{DNN}}^* = \min \text{ trace } L_s Y + \mathbb{1}_{\mathcal{Y}}(Y) + \mathbb{1}_{\mathcal{R}}(R) \tag{3.3}$$

$$\text{s.t. } Y = \widehat{V}R\widehat{V}^T,$$

where

$$\mathcal{R} := \left\{ R \in \mathbb{S}_+^{(k-1)(n-1)+1} : \text{trace } R = n + 1 \right\};$$

$$\mathcal{Y} := \{Y \in \mathbb{S}^{nk+1} : \mathcal{G}_{\widehat{J}_0}(Y) = \mathcal{G}_{\widehat{J}_0}(u_0 u_0^T), 0 \leq \mathcal{G}_{\widehat{J}_0}(Y) \leq 1,$$

$$D_o(Y) = \widehat{M}, e^T Y_{(i0)} = m_i, i = 1, \dots, k\}.$$

Notice that the sets  $\mathcal{R}, \mathcal{Y}$  are much smaller than  $\mathcal{R}_o, \mathcal{Y}_o$ , respectively. This property can help bring the iterates of  $R, Y$  closer to the optimal solution set more quickly,



when a splitting method is applied. In addition, we see below in Sects. 3.1.1 and 3.1.2, that these redundant constraints do not significantly increase the computational cost.

We now describe our splitting method for solving (3.3) (which is equivalent to solving (2.50)). We start by writing down the augmented Lagrangian function for (3.3):

$$\mathcal{L}_\beta(R, Y, Z) = f_{\mathcal{R}}(R) + g_{\mathcal{Y}}(Y) + \langle Z, Y - \widehat{V}R\widehat{V}^T \rangle + \frac{\beta}{2} \|Y - \widehat{V}R\widehat{V}^T\|^2,$$

where  $\beta > 0$  is a penalty parameter for the quadratic penalty term, and  $f_{\mathcal{R}}(R)$  and  $g_{\mathcal{Y}}(Y)$  are defined respectively as

$$f_{\mathcal{R}}(R) = \mathbb{1}_{\mathcal{R}}(R), \quad g_{\mathcal{Y}}(Y) = \text{trace } L_s Y + \mathbb{1}_{\mathcal{Y}}(Y).$$

Our main Algorithm 3.1 for solving (3.3) is a standard application of the *strictly contractive Peaceman-Rachford splitting method, sPRSM* [11] to (3.3). It can be summarized as follows:

- alternate minimization of  $\mathcal{L}_\beta$  in the variables  $Y$  and  $R$  interlaced by an update of the  $Z$  variable;
- update the dual variable  $Z$  both after the  $R$ -update *and* the  $Y$ -update (both  $R$  and  $Y$  updates in (3.4) are well defined as both constraint sets are closed, convex, and  $\widehat{V}^T \widehat{V} = I$ ).

---

**Algorithm 3.1: FRSMR for DNN relaxation**

---

**Step 1.** Pick any  $Y^0, Z^0 \in \mathbb{S}^{nk+1}$ . Fix  $\beta > 0$  and  $\gamma \in (0, 1)$ . Set  $t = 0$ .

**Step 2.** For each  $t = 0, 1, \dots$ , update (precise stopping criteria: Section 4.2, item 2)

$$\begin{aligned} R^{t+1} &= \arg \min_{R \in \mathcal{R}} \mathcal{L}_\beta(R, Y^t, Z^t) = \arg \min_R f_{\mathcal{R}}(R) - \langle Z^t, \widehat{V}R\widehat{V}^T \rangle + \frac{\beta}{2} \|Y^t - \widehat{V}R\widehat{V}^T\|^2, \\ Z^{t+\frac{1}{2}} &= Z^t + \gamma\beta(Y^t - \widehat{V}R^{t+1}\widehat{V}^T), \\ Y^{t+1} &= \arg \min_{Y \in \mathcal{Y}} \mathcal{L}_\beta(R^{t+1}, Y, Z^{t+\frac{1}{2}}) = \arg \min_Y g_{\mathcal{Y}}(Y) + \langle Z^{t+\frac{1}{2}}, Y \rangle + \frac{\beta}{2} \|Y - \widehat{V}R^{t+1}\widehat{V}^T\|^2, \\ Z^{t+1} &= Z^{t+\frac{1}{2}} + \gamma\beta(Y^{t+1} - \widehat{V}R^{t+1}\widehat{V}^T). \end{aligned} \tag{3.4}$$


---

We next discuss convergence of the sequence generated by Algorithm 3.1. Recall from Proposition 2.17 that (2.50) has primal Slater points. Consequently,  $(Y^*, R^*)$  solves (3.3) if, and only if, there exists  $Z^*$  so that the following first order optimality conditions hold:

$$\begin{aligned} 0 &\in -\widehat{V}^T Z^* \widehat{V} + \mathcal{N}_{\mathcal{R}}(R^*), \\ 0 &\in L_s + Z^* + \mathcal{N}_{\mathcal{Y}}(Y^*), \\ Y^* &= \widehat{V}R^* \widehat{V}^T, \end{aligned} \tag{3.5}$$

where  $\mathcal{N}_S(x)$  denotes the normal cone of  $S$  at  $x$ . The following Theorem 3.1 states that the sequence generated by Algorithm 3.1 converges to a point satisfying (3.5). Its proof can be found in [11].

**Theorem 3.1** *Let  $\{R^t\}, \{Y^t\}, \{Z^t\}$  be the sequences generated by Algorithm 3.1. Then  $\{(R^t, Y^t)\}$  converges to an optimal solution  $(R^*, Y^*)$  of (3.3), and  $\{Z^t\}$  converges to some  $Z^*$  so that  $(R^*, Y^*, Z^*)$  satisfies (3.5).*

In Algorithm 3.1, the explicit  $Z$ -update in (3.4) is simple and easy. We now show that we have explicit expressions for the  $R, Y$ -updates as well.

### 3.1.1 R-subproblem

Recall that Assumption 2.15 guarantees that  $\widehat{V}$  is normalized so that  $\widehat{V}^T \widehat{V} = I$ . Then the  $R$ -subproblem can be explicitly solved by projecting onto the set  $\mathcal{R}$

$$\begin{aligned} R^{t+1} &= \arg \min_{R \in \mathcal{R}} -\langle Z^t, \widehat{V}R\widehat{V}^T \rangle + \frac{\beta}{2} \left\| Y^t - \widehat{V}R\widehat{V}^T \right\|^2 \\ &= \arg \min_{R \in \mathcal{R}} \frac{\beta}{2} \left\| Y^t - \widehat{V}R\widehat{V}^T + \frac{1}{\beta} Z^t \right\|^2 \\ &= \arg \min_{R \in \mathcal{R}} \frac{\beta}{2} \left\| R - \widehat{V}^T (Y^t + \frac{1}{\beta} Z^t) \widehat{V} \right\|^2 \\ &= \mathcal{P}_{\mathcal{R}} (\widehat{V}^T (Y^t + \frac{1}{\beta} Z^t) \widehat{V}), \end{aligned}$$

where  $\mathcal{P}_{\mathcal{R}}$  denotes the projection (nearest point) onto the intersection of the positive semidefinite cone  $\mathbb{S}_+^{(k-1)(n-1)+1}$  and the hyperplane  $\{R \in \mathbb{S}^{(k-1)(n-1)+1} : \text{trace } R = n + 1\}$ . For any symmetric matrix  $W \in \mathbb{S}^{(n-1)(k-1)+1}$ , we have

$$\mathcal{P}_{\mathcal{R}}(W) = U \text{diag}(\mathcal{P}_{\bar{\Lambda}}(\text{diag}(\Lambda))) U^T,$$

where  $(U, \Lambda)$  contains the eigenpairs of  $W$ , and  $\mathcal{P}_{\bar{\Lambda}}$  denotes the projection of the vector of eigenvalues, i.e.,  $\text{diag}(\Lambda)$ , onto the simplex  $\bar{\Lambda} = \{\lambda \in \mathbb{R}_+^{(k-1)(n-1)+1} : \lambda^T e = n + 1\}$ . Projection onto simplices can be performed efficiently via some standard root-finding strategies; see, for example, [6, 28].

### 3.1.2 Y-subproblem

The  $Y$ -subproblem involves projection onto the polyhedral set  $\mathcal{Y}$ , i.e.,

$$\begin{aligned} Y^{t+1} &= \arg \min_{Y \in \mathcal{Y}} \langle L_s, Y \rangle + \langle Z^{t+\frac{1}{2}}, Y - \widehat{V}R^{t+1}\widehat{V}^T \rangle + \frac{\beta}{2} \left\| Y - \widehat{V}R^{t+1}\widehat{V}^T \right\|^2 \\ &= \arg \min_{Y \in \mathcal{Y}} \frac{\beta}{2} \left\| Y - \widehat{V}R^{t+1}\widehat{V}^T + \frac{1}{\beta} (L_s + Z^{t+\frac{1}{2}}) \right\|^2. \end{aligned} \tag{3.6}$$

To present a closed form solution for the update, we let  $Y := \widehat{V}R^{t+1}\widehat{V}^T - \frac{1}{\beta}(L_s + Z^{t+\frac{1}{2}})$  and assume that  $Y$  is blocked as in (1.4). We now partition the set of indices of  $J_0^c$  into the following three disjoint sets:

- $\zeta_r$ : it includes the 0-th row of  $Y$  except for the 00-element.
- $\zeta_o(\subseteq J_0^c)$ : it includes all off-diagonal elements of the blocks in  $Y$  whenever these off-diagonal elements belong to  $J_0^c$ .
- $\zeta_d$ : it includes the diagonal of  $Y$  except for the 00-element.

We also define the following subsets:

$$\begin{aligned} \mathcal{Y}_g &:= \{Y \in \mathbb{S}^{nk+1} : \mathcal{G}_{J_0}(Y) = \mathcal{G}_{J_0}(u_0 u_0^T)\}; \\ \mathcal{Y}_r &:= \{Y \in \mathbb{S}^{nk+1} : 0 \leq \mathcal{G}_{\zeta_r}(Y) \leq 1, e^T Y_{(i0)} = m_i, i = 1, \dots, k\}; \\ \mathcal{Y}_o &:= \{Y \in \mathbb{S}^{nk+1} : 0 \leq \mathcal{G}_{\zeta_o}(Y) \leq 1, \mathcal{D}_o(Y) = \widehat{M}\}; \\ \mathcal{Y}_d &:= \{Y \in \mathbb{S}^{nk+1} : 0 \leq \mathcal{G}_{\zeta_d}(Y) \leq 1\}. \end{aligned}$$

Note that  $\mathcal{Y} = \mathcal{Y}_g \cap \mathcal{Y}_d \cap \mathcal{Y}_r \cap \mathcal{Y}_o$ . The next iterate  $Y^{t+1}$  can now be computed as follows:

$$(Y^{t+1})_{ij} = \begin{cases} 1 & \text{if } i = j = 0, \\ 0 & \text{if } ij \in J_0, \\ (\mathcal{P}_{\mathcal{Y}_r}(Y))_{ij} & \text{if } ij \in \zeta_r, \\ (\mathcal{P}_{\mathcal{Y}_o}(Y))_{ij} & \text{if } ij \in \zeta_o, \\ \min(1, \max(Y_{ij}, 0)) & \text{if } ij \in \zeta_d, \end{cases}$$

where  $\mathcal{P}_{\mathcal{Y}_r}$  and  $\mathcal{P}_{\mathcal{Y}_o}$  denote the orthogonal projection onto the  $\mathcal{Y}_r$  and  $\mathcal{Y}_o$  respectively. Both  $\mathcal{Y}_r$  and  $\mathcal{Y}_o$  are intersections of a hyperplane and a box. The projection can be obtained efficiently via standard root-finding algorithms; see, for example, [14, 17].

Denote the inexact approximate solution from **FRSMR** by  $(R^{\text{out}}, Y^{\text{out}}, Z^{\text{out}})$ . In the following two subsections, we illustrate how we compute the lower and upper bounds with the obtained  $Z^{\text{out}}$  and  $Y^{\text{out}}$ , respectively.

### 3.2 Lower bound from inaccurate relaxation

Since (3.3) is a relaxation of **MC**, we conclude that exact solutions provide a lower bound for the original **MC**. However, the problem size of (3.3) can be extremely large, and it is generally too expensive to obtain highly accurate solutions. In the following, we provide an inexpensive way to get a valid lower bound from the output of our algorithm even when the solution is only obtained to a moderate accuracy. Our approach is based on the following function

$$g(Z) := \min_{Y \in \widehat{\mathcal{Y}}} \langle L_s + Z, Y \rangle - (n + 1)\lambda_{\max}(\widehat{V}^T Z \widehat{V}), \tag{3.7}$$

where  $\lambda_{\max}(\widehat{V}^T Z \widehat{V})$  denotes the largest eigenvalue of  $\widehat{V}^T Z \widehat{V}$ , and the constraint set

$$\begin{aligned}\tilde{\mathcal{Y}} &:= \{Y \in \mathbb{S}^{nk+1} : \mathcal{G}_{j_0}(Y) = \mathcal{G}_{j_0}(u_0 u_0^T), 0 \leq \mathcal{G}_{j_0}(Y) \leq 1, \\ &\quad \mathcal{D}_o(Y) = \hat{M}, \mathcal{D}_i(Y) = M, e^T Y_{(i0)} = m_i, i = 1, \dots, k\}.\end{aligned}$$

In the following Theorem 3.2, we show that  $\max_Z g(Z)$  is indeed a Fenchel dual problem of (3.3). Since the Fenchel dual problem is an unconstrained maximization problem, evaluating  $g$  in (3.7) at the  $t$ -th iterate  $Z^t$  returned by Algorithm 3.1 always yields a lower bound for  $p_{\text{DNN}}^*$ .<sup>7</sup>

**Theorem 3.2** Consider the problem

$$d_Z^* := \max_Z g(Z), \quad (3.8)$$

where  $g$  is defined in (3.7). Then (3.8) is a concave maximization problem and strong duality holds between (3.3) and (3.8), i.e.,

$$d_Z^* = p_{\text{DNN}}^*, \text{ and } d_Z^* \text{ is attained.}$$

**Proof** We derive (3.8) as a Fenchel dual problem of (3.3) by finding a best lower bound as follows.

$$\begin{aligned}p_{\text{DNN}}^* &= \min_{R \in \mathcal{R}, Y \in \mathcal{Y}} \max_Z \left\{ \langle L_s, Y \rangle + \langle Z, Y - \hat{V}R\hat{V}^T \rangle \right\} \\ &= \min_{R \in \mathcal{R}, Y \in \tilde{\mathcal{Y}}} \max_Z \left\{ \langle L_s, Y \rangle + \langle Z, Y - \hat{V}R\hat{V}^T \rangle \right\}\end{aligned} \quad (3.9a)$$

$$= \max_Z \min_{R \in \mathcal{R}, Y \in \tilde{\mathcal{Y}}} \left\{ \langle L_s, Y \rangle + \langle Z, Y - \hat{V}R\hat{V}^T \rangle \right\} \quad (3.9b)$$

$$\begin{aligned}&= \max_Z \left\{ \min_{Y \in \tilde{\mathcal{Y}}} \left\{ \langle L_s, Y \rangle + \langle Z, Y \rangle \right\} + \min_{R \in \mathcal{R}} \langle Z, -\hat{V}R\hat{V}^T \rangle \right\} \\ &= \max_Z \left\{ \min_{Y \in \tilde{\mathcal{Y}}} \left\{ \langle L_s, Y \rangle + \langle Z, Y \rangle \right\} + \min_{R \in \mathcal{R}} \langle \hat{V}^T Z \hat{V}, -R \rangle \right\}\end{aligned} \quad (3.9c)$$

$$= \max_Z \left\{ \min_{Y \in \tilde{\mathcal{Y}}} \langle L_s + Z, Y \rangle - (n+1)\lambda_{\max}(\hat{V}^T Z \hat{V}) \right\} = d_Z^*, \quad (3.9d)$$

where:

- (3.9a) follows from the redundancy of the constraint  $\mathcal{D}_i(Y) = M$  as guaranteed by Theorem 2.11;<sup>8</sup>

<sup>7</sup> This strengthens [18, Lemma 3.2].

<sup>8</sup> Note that the inner maximization forces  $Y = \hat{V}R\hat{V}^T$ .

2. (3.9b) follows from [24, Corollary 28.2.2], [24, Theorem 28.4] and the fact that (3.3) has Slater points (see Proposition 2.17);<sup>9</sup>
3. (3.9d) follows from the definition of  $\mathcal{R}$  and the Rayleigh Principle.

The concavity of  $g$  is clear, and we see from [24, Corollary 28.2.2] and [24, Corollary 28.4.1] that the dual value  $d_Z^*$  is attained. □

### 3.3 Upper bound from a feasible solution

We now move from lower bounds to finding upper bounds for cut ( $m$ ). Let  $Y^{\text{out}}$  be the output from our algorithm **FRSMR**, i.e.,  $Y^{\text{out}}$  is obtained from solving (3.6). The [procedures for computing upper bounds are:]

1. We extract a column vector  $v$  from  $Y^{\text{out}}$  in one of the following three ways:<sup>10</sup>
  - (a) use column 0 of  $Y^{\text{out}}$ ;
  - (b) use the eigenvector corresponding to the largest eigenvalue of  $Y^{\text{out}}$ ;
  - (c) sum of random weighted-eigenvalue eigenvectors of  $Y^{\text{out}}$ , i.e.,

$$v = \sum_{i=1}^r w_i \lambda_i v_i,$$

where  $\lambda_1 \geq \dots \geq \lambda_r > 0$ , are the ordered eigenvalues of  $Y^{\text{out}}$  with eigenpairs  $(\lambda_i, v_i)$ , and  $1 \geq w_1 \geq \dots \geq w_r > 0$  are random ordered weights. The  $r$  here is the *numerical rank* of  $Y^{\text{out}}$ .<sup>11</sup>

2. For each vector  $v$  obtained in Step 1, we extract its last  $nk$  elements as a subvector  $v^\circ$  and set  $X^\circ = \text{Mat}(v^\circ)$ .
3. For each  $X^\circ$  obtained, we find the nearest partition matrix  $X^*$  to it. (See Proposition 3.4, below.)
4. For each  $X^*$  obtained, an upper bound of **MC** is found as  $\frac{1}{2} \text{trace}(AX^*BX^{*T})$ . We save the best (smallest) upper bound obtained and the corresponding  $X^*$ . (We repeat the random choice in Item 1c  $\lceil \log(n) \rceil$  times.)

<sup>9</sup> Note that the Lagrangian is linear in  $R, Y$  and linear in  $Z$ . Moreover, both constraint sets  $\mathcal{R}, \mathcal{Y}$  are convex and compact. Therefore, the result also follows from the classical Von Neumann-Fan minmax theorem.

<sup>10</sup> Note that if  $Y^{\text{out}}$  is rank-1 and feasible, then the first two methods in Item 1a and Item 1b yield exact solutions to **MC**. This motivates the use of eigenvector information.

<sup>11</sup> MATLAB:  $r = \min(\text{sum}(\lambda/(n+1) > 0.1) + 1, n + 1)$ ;

**Remark 3.3**

1. First of all, the projection in Item 3 can be done efficiently using linear programming. (Actually in strongly polynomial time if one uses something like the classical Hungarian algorithm.) This is similar to what is done in [18, 19, 31].
2. In [18], we adopt a similar procedure for calculating an upper bound but with only one column vector  $v$  from  $Y^{\text{out}}$  using Items 1a and 1b. In Figure 1, we compare with the method in [18] and see that Item 1c provides a significant improvement to the upper bound.

**Proposition 3.4** [19, Theorem 6.1] *Let  $X^\circ \in \mathbb{R}^{n \times k}$ . Then the nearest partition matrix  $X^* \in \mathcal{M}_m$  to  $X^\circ$  can be found by solving the transportation type linear program*

$$\begin{aligned} X^* \in \arg \min \quad & - \text{trace } X^\circ T X \\ \text{s.t.} \quad & X e = e \\ & X^T e = m \\ & X \geq 0. \end{aligned} \quad (3.10)$$

Note that we get an exact solution if  $\text{rank}(Y^{\text{out}}) = 1$  and  $Y^{\text{out}} = \widehat{V} R^{\text{out}} \widehat{V}^T$ . Lemma 3.5 below suggests that the methods described in Item 1a and Item 1b above likely yield reasonable approximate partition matrices. Recall that

$$\text{conv } \mathcal{M}_m = \{X \in \mathbb{R}^{n \times k} : X e = e, X^T e = m, X \geq 0\}.$$

**Proposition 3.5** [19, Proposition 5.2] *Let  $Y$  be feasible for (2.50). Let  $v_1 = Y_{1:nk0}$ , and let  $[v_0 \ v_2^T]^T$  denote a unit eigenvector of  $Y$  corresponding to the largest eigenvalue. Then  $v_0 \neq 0$ , and both*

$$X_1^\circ := \text{Mat}(v_1), X_2^\circ := \text{Mat}(v_0^{-1} v_2) \in \text{conv } \mathcal{M}_m.$$

However, in general  $Y^{\text{out}}$  is not an exact solution of the DNN relaxation. Then Item 1c in Remark 3.3 plays an important role in generating many vectors  $v$  for finding an upper bound. We see this in Section 4.3.3 below. In fact, this allows us to stop the algorithm with much fewer iterations when we see that both the upper and lower bounds are not improving.

## 4 Numerical experiments

In this section we apply the proposed **FRSMR** method in Algorithm 3.1 to solve the **DNN** relaxation in (3.3). All the tests are performed using Matlab R2017a on a ThinkPad X1 with an Intel CPU (2.5GHz) and 8GB RAM running Windows 10.

### 4.1 Classes of problems and parameters

We consider three classes of problems, see Sections 4.3.1 to 4.3.3. We outline them here:

- (a) (random structured graphs, Section 4.3.1.) We compare with the **DNN** relaxation in [19].<sup>12</sup> The latter relaxation is solved using an interior point approach with Mosek version 8.0.0.60. [1]. See Table 2.
- (b) (partially random graphs with various sizes, Section 4.3.2.) There are four kinds of random graphs, classified by the number of 1's,  $|\mathcal{I}|$ , in the vector  $m$ . In particular, in the three cases where  $\mathcal{I} \neq \emptyset$ , we *almost always* obtain a zero gap and thus the optimal solution. See Tables 3, 4, 5 and 6.
- (c) (vertex separator instances, Section 4.3.3.) We compare with the bounds obtained by solving the relaxation **SDP**<sub>4</sub> in [22]. In addition, we include comparisons on the upper bounds on the size of the vertex separator. See Table 7.

## 4.2 Parameters, initialization, stopping criteria

In our implementation, we first shift the objective to obtain positive definiteness.

$$L \leftarrow L + \alpha I, \quad \alpha = 0.1 + \max\{0, -\lambda_{\min}(L)\}.$$

This does not change the optimum  $Y^*$  but it changes the dual  $Z$  and promotes  $Z \leq 0$ , as can be seen from the expression for the  $Y$ -subproblem in (3.6). This in turn promotes a better lower bound from (3.9d).

We now specify the parameters used in **FRSMR** in Sections 4.3.1 to 4.3.3.

1. The penalty and step parameters are  $\beta = 3k/n$  and  $\gamma = 0.9$ , respectively.
2. We terminate once one of the following Items 2a to 2c holds:
  - (a) the number of iterations reaches 10000;
  - (b) the relative gap, rel-gap, is either zero<sup>13</sup> or does not change in  $\max\{5, \lceil n/10 \rceil\}$  consecutive iterations,
  - (c)  $\text{rel-gap} = \frac{(\text{best upper bound} - \text{best lower bound})}{(\text{best upper bound} + \text{best lower bound} + 1)/2}$ ;

$$\max \left\{ \left\| Y^{t+1} - \widehat{V} R^{t+1} \widehat{V}^T \right\|, \left\| Y^{t+1} - Y^t \right\| \right\} < 10^{-12}; \tag{3.1}$$

This criterion (3.1) is the same as that suggested in [12, Remark 2.3].

3. We calculate: the lower bound and the upper bound every 100th iteration, using Theorem 3.2 (to compute a lower bound as  $\lceil g(Z^t) \rceil$ ) and the procedures in Section 3.3. In the computation of the upper bound, we sample the random weight vector  $\lceil \log(n) \rceil$  times. The linear program (3.10) involved in the computation of

<sup>12</sup> The **DNN** relaxation in [19] imposes the additional nonnegativity constraints  $\widehat{V} Z \widehat{V}^T \geq 0$  onto their **SDP**<sub>final</sub> relaxation.

<sup>13</sup> Note that our data are integral and we round up the lower bound, therefore the gap is integer valued. Thus, finding a zero duality gap is reasonable. Moreover, the lower bounds are nonnegative.

**Table 1** Data terminology

imax	The maximum size of each set
$k$	The number of sets
$n$	The number of nodes, i.e., the sum of the sizes of the sets
$p$	The density of the graph, i.e., $2 E /( V ( V  - 1))$
$l = e^T m_{\text{one}}$	The number of 1's in $m$
Iters	The number of iterations
Time	CPU time in seconds
Bounds	Best lower and upper bounds and relative gap
Residuals	Final values $\ Y^{r+1} - \hat{V}R^{r+1}\hat{V}^T\ $ ( $\cong \Delta Z$ ); $\ Y^{r+1} - Y^r\ $ ( $\cong \Delta Y$ )

**Table 2** Small structured graphs; comparisons with DNN relaxation model in [19]

Data				Lower bounds		Upper bounds		Rel-gap		Time (cpu)	
$n$	$k$	$ E $	$u_0$	<b>FRSMR</b>	Mosek	<b>FRSMR</b>	Mosek	<b>FRSMR</b>	Mosek	<b>FRSMR</b>	Mosek
20	4	136	6	6	6	6	6	0.00	0.00	0.14	5.41
25	4	222	8	8	8	8	8	0.00	0.00	0.22	10.24
25	5	170	14	14	14	14	14	0.00	0.00	0.27	30.36
31	5	265	22	22	22	22	22	0.00	0.00	1.15	126.11

the upper bound is solved with Mosek using their function ‘mosekopt’ and the dual-simplex method.

4. The data terminology in our Tables are described in Table 1.
5. In Sect. 4.3.3 we consider the special class of vertex separator problems.
  - (a) The penalty and step parameters in **FRSMR** are  $\beta = 0.001$  and  $\gamma = 0.9$  respectively.
  - (b) The stopping criterion is set as the same as in Sects. 4.3.1 and 4.3.2.
  - (c) We calculate the lower bound every 100-th iteration using Theorem 3.2. We compute the upper bound every iteration using the procedures in Sect. 3.3. Other settings in the computation of the upper bound are the same as in Sects. 4.3.1 and 4.3.2.

### 4.3 Three classes of problems

#### 4.3.1 Random structured graphs

The structured graphs are generated as in [19, Sect. 7.1]. That is, we first generate  $k$  disjoint cliques of sizes  $m_1, \dots, m_k$ , randomly chosen from  $\{2, \dots, \text{imax}\}$ . We then join the first  $k - 1$  cliques to every node of the  $k$ -th clique, and add  $u_0$  edges between the first  $k - 1$  cliques, chosen uniformly at random from the complement graph. In



**Table 3** Results for random graphs with  $\mathcal{I} = \emptyset$

Specifications					Iters	Time (cpu)	Bounds			Residuals	
imax	$k$	$n$	$p$	$l$			Lower	Upper	Rel-gap	Primal	Dual
4	5	17	0.43	0	500	0.94	16	17	0.06	9.51e-04	1.01e-04
4	5	17	0.32	0	100	0.19	10	10	0.00	1.93e-02	1.75e-02
5	6	23	0.35	0	500	1.75	37	42	0.13	1.81e-03	1.92e-04
5	6	23	0.30	0	600	1.92	30	34	0.12	1.07e-03	1.68e-04
6	7	30	0.28	0	900	5.99	42	48	0.13	1.65e-03	1.28e-04
6	7	30	0.22	0	600	4.14	31	40	0.25	3.24e-03	3.88e-04
7	8	37	0.18	0	700	9.03	32	38	0.17	6.29e-03	1.56e-03
7	8	37	0.14	0	700	9.13	18	22	0.20	5.22e-03	1.18e-03
8	9	49	0.10	0	1200	47.09	14	19	0.29	5.68e-03	8.18e-04
8	9	49	0.05	0	1000	45.52	0	6	1.71	1.31e-04	1.83e-04

our experiments below, we set  $u_0 = \lfloor e_c d \rfloor$ , where  $e_c$  is the number of edges in the complement graph and  $d$  is the density (percentage of edges in the complement graph to be added). By construction,  $u_0 \geq \text{cut}(m)$ .

We use small instances with  $k = 4, 5, d = 10\%$  and  $\text{imax} = 6, 8$ . We compare our approach with the DNN relaxation model in [19] solved by Mosek [1]. The results in Table 2 illustrate the improvement in solution time.

### 4.3.2 (Partially) random graphs with various sizes

We test four groups of random graphs corresponding to different values of  $\mathcal{I}$ :

1. ( $\mathcal{I} = \emptyset$ ) vector  $m$  is generated by choosing  $k$  integers randomly from  $\{2, \dots, \text{imax}\}$  ;
2. ( $k \notin \mathcal{I} \neq \emptyset$ ) after generating  $m$  as in Item 1 above, we randomly select elements from  $\{m_1, m_2, \dots, m_{k-1}\}$  and set them to be 1;<sup>14</sup>
3. ( $k \in \mathcal{I} \neq \mathcal{K}$ ) after generating  $m$  as in Item 1 above, we set  $m_k = 1$  and randomly select no more than  $k - 2$  elements from  $\{m_1, m_2, \dots, m_{k-1}\}$  and set them to be 1;
4. ( $\mathcal{I} = \mathcal{K}$ ) simply set  $\text{imax} = 1$  and set all the elements of  $m$  to be 1.<sup>15</sup>

<sup>14</sup> In this case and the next, we have  $m_i = 1$  for some  $i$ . One typical application is the side chain positioning problem that involves rotamer selections in protein folding. Some rotamer sets typically often only one rotamer, see [5, 13].

<sup>15</sup> As pointed out by one reviewer, in this case, we have  $n = k$ , and the objective function in (1.1) is the number of nonzero entries that remain in the upper triangular part of  $X^TAX$  after removing the last row and column. Thus, an optimal solution  $X \in \mathcal{M}_e$  is a permutation matrix that puts the vertex with the highest degree as vertex  $n$ . Here, we include these instances to test the effectiveness of our new gangster constraints.

**Table 4** Results for random graphs with  $k \notin \mathcal{I} \neq \emptyset$

Specifications					Iters	Time (cpu)	Bounds			Residuals	
imax	$k$	$n$	$p$	$l$			Lower	Upper	Rel-gap	Primal	Dual
4	5	14	0.37	1	100	0.17	6	6	0.00	1.59e-02	1.26e-02
4	5	14	0.37	1	100	0.17	5	5	0.00	2.88e-02	4.62e-02
5	6	16	0.35	2	400	0.92	11	11	0.00	1.70e-03	4.32e-04
5	6	16	0.32	2	100	0.24	11	11	0.00	2.81e-02	3.22e-02
6	7	19	0.27	4	500	1.79	8	9	0.11	2.73e-03	3.29e-04
6	7	19	0.22	4	500	1.76	4	5	0.20	1.75e-03	4.32e-04
7	8	12	0.20	7	100	0.21	0	0	0.00	1.20e-02	1.54e-02
7	8	12	0.17	7	100	0.21	0	0	0.00	2.19e-02	1.97e-02
8	9	16	0.12	8	100	0.38	0	0	0.00	4.78e-02	6.50e-02
8	9	16	0.06	8	100	0.38	0	0	0.00	3.06e-02	3.10e-02

**Table 5** Results for random graphs with  $k \in \mathcal{I} \neq \mathcal{K}$

Specifications					Iters	Time (cpu)	Bounds			Residuals	
imax	$k$	$n$	$p$	$l$			Lower	Upper	Rel-gap	Primal	Dual
4	5	12	0.45	2	100	0.16	11	11	0.00	1.41e-03	2.03e-03
4	5	12	0.39	2	100	0.14	9	9	0.00	1.08e-02	1.38e-02
5	6	15	0.33	3	100	0.21	13	13	0.00	2.43e-02	3.80e-02
5	6	15	0.29	3	100	0.21	10	10	0.00	3.12e-02	5.09e-02
6	7	18	0.27	4	100	0.37	13	13	0.00	8.97e-02	1.03e-01
6	7	18	0.22	4	300	0.95	10	10	0.00	3.82e-03	2.76e-03
7	8	13	0.21	7	100	0.23	5	5	0.00	7.67e-03	8.75e-03
7	8	13	0.18	7	100	0.23	4	4	0.00	1.56e-02	1.94e-02
8	9	16	0.11	8	100	0.47	2	2	0.00	5.51e-02	1.04e-01
8	9	16	0.06	8	100	0.49	0	0	0.00	1.30e-02	1.47e-02

Then, as  $n = m^T e$  is the total number of nodes in the simple, undirected graph, we randomly generate an adjacency matrix  $A$  of a graph on  $n$  nodes with density =  $\text{density}A$ , and construct the Laplacian matrix.<sup>16</sup>

In Tables 3 to 6, we consider the four groups of random graphs in Items 1 to 4, above. In each group of random graphs, we generate  $m$  and  $A$  by choosing  $k$  and imax as given in the tables with various values for  $\text{density}A$ ; the density  $p$  of the graphs is also reported.

<sup>16</sup> In MATLAB:  $A = \text{abs}(\text{sprandsym}(\text{sum}(m), \text{density}A)) > 0; A = A - \text{diag}(\text{diag}(A));$  Note that  $\text{density}A$  is different from the density of the graph  $p$  defined in Table 1.

**Table 6** Results for random graphs with  $\mathcal{I} = \mathcal{K}$

Specifications					Iters	Time (cpu)	Bounds			Residuals	
imax	k	n	p	l			Lower	Upper	Rel-gap	Primal	Dual
1	8	8	0.64	8	100	0.17	12	12	0.00	4.22e-04	6.08e-04
1	10	10	0.69	10	100	0.26	23	23	0.00	9.94e-03	1.26e-02
1	12	12	0.47	12	100	0.39	23	23	0.00	1.86e-02	3.32e-02
1	14	14	0.46	14	100	0.66	33	33	0.00	6.37e-02	8.99e-02
1	16	16	0.44	16	100	1.04	43	43	0.00	1.69e-01	2.49e-01
1	18	18	0.39	18	200	3.71	48	48	0.00	1.45e-02	2.22e-02
1	20	20	0.29	20	200	7.31	47	47	0.00	3.75e-02	4.04e-02
1	22	22	0.25	22	200	11.24	47	47	0.00	1.39e-01	1.58e-01
1	24	24	0.13	24	200	16.41	31	31	0.00	1.06e-01	1.13e-01
1	26	26	0.05	26	200	23.75	10	10	0.00	1.19e-01	8.14e-02

From Table 3, i.e., in the case of  $\mathcal{I} = \emptyset$ , we can see that the **FRSMR** in general takes a reasonably short time to converge. Moreover, in most instances, the rel-gap is very small; sometimes we even obtain a zero gap and hence the instance is solved to optimality. **FRSMR** appears to perform better in the cases when  $\mathcal{I} \neq \emptyset$ . The corresponding results are shown in Tables 4 to 6. We can see that in most instances, the rel-gap is zero and the problem is solved exactly. Moreover, the CPU times taken are reasonably small.

### 4.3.3 Vertex separator problem

We now test some vertex separator problems from <https://sites.google.com/site/sotirovr/the-vertex-separator>. We compare against the bounds obtained from the model **SDP**<sub>4</sub> in [22]. In each instance, the  $m$  has the special structure that  $k = 3$ ,  $|m_1 - m_2| \leq 1$  and  $\text{cut}(m) > 0$ . In this case, by solving **MC**, one can separate the nodes of the graph into  $S_1, S_2$  and  $S_3$  so that the number of edges between  $S_1$  and  $S_2$  is minimized. If  $\text{cut}(m) = 0$ , for some  $m = (m_1, m_2, m_3)^T$ , then we say that  $S_3$  separates  $S_1$  and  $S_2$ , and  $S_3$  is called a *vertex separator*. If  $\text{cut}(m) > 0$ , on the other hand, it means that no separator  $S_3$  for the cardinalities specified in  $m$  exists. However, we can experiment with different choices of  $m$ , i.e., transferring nodes from  $S_1$  and  $S_2$  to  $S_3$ , in the hope of eventually producing a separator. In this way, we can obtain an upper bound of the cardinality of a vertex separator. Here, we follow the approach described in [22, Section 8] to derive an upper bound of the cardinality of a vertex separator, using solutions obtained from **FRSMR**.

In Table 7, we compare the lower and upper bounds for  $\text{cut}(m)$  obtained from (3.3) and from the model **SDP**<sub>4</sub> in [22]. We also report the upper bound of the cardinality of vertex separator obtained for each instance. The (upper and lower)

**Table 7** Comparing bounds for MC and bounds for the cardinality of separators

Name	$n$	$ E $	$m_1$	$m_2$	$m_3$	MC by $SDP_4$		MC by (3.3)		Separator by $SDP_4$		Separator by (3.3)	
						Lower	Upper	Lower	Upper	Lower	Upper	Lower	Upper
Example 1	93	470	42	41	10	0.07	1	0	1	11	11	11	11
bcspr03	118	179	58	57	3	0.56	1	0	2	4	5	5	5
Smallmesh	136	354	65	66	5	0.13	1	0	1	6	6	6	6
can-144	144	576	70	70	4	0.90	6	0	6	5	6	8	8
can-161	161	608	73	72	16	0.31	2	0	2	17	18	18	18
can-229	229	774	107	107	15	0.40	6	0	6	16	19	19	19
gridt(15)	120	315	56	56	8	0.29	4	0	4	9	11	12	12
gridt(17)	153	408	72	72	9	0.17	4	0	4	10	13	13	13
grid3dt(5)	125	604	54	53	18	0.54	2	0	4	19	19	22	22
grid3dt(6)	216	1115	95	95	26	0.28	4	0	4	27	30	31	31
grid3dt(7)	343	1854	159	158	26	0.60	22	0	27	27	37	44	44




Fig. 1 Comparison of upper bounds with/without randomness; Item 1c, Sect. 3.3

bounds for  $SDP_4$  are obtained directly from [22, Table 3].<sup>17</sup> From Table 7, we can see that the MC upper bounds from the model (3.3) are very competitive with those obtained from the model  $SDP_4$ . For most instances, the upper bounds are equal except for two instances, “grid3dt(5)” and “grid3dt(7)”; as for the comparison of upper bounds for vertex separator, still most upper bounds are equal, except for “can-144”, “gridt(15)”, “gridt(5)”, “gridt(6)” and “gridt(7)”.

Figure 1 shows the comparison between the upper bound using Items 1a to 1c in Sect. 3.3 and the one using only Items 1a and 1b there. It shows that the former strategy can produce much better upper bounds than those obtained by the latter strategy.

## 5 Conclusion

In this paper we introduced new methods for finding strengthened lower and upper bounds for the MC problem. SDP relaxations provide strong bounds that are further strengthened by nonnegativity constraints, i.e., by using the DNN relaxation. However, in general solving the DNN relaxation by interior-point methods is extremely expensive.

The FR appears to provide a natural splitting for the variables  $Y = \widehat{V}R\widehat{V}^T$ , where  $Y, R$  are restricted to the polyhedral and cone constraints, respectively. We exploit this within a sPRSM framework.

<sup>17</sup> These results use extra cutting planes, and therefore they obtain stronger lower bounds on cut ( $m$ ).

We bring back previously redundant constraints to strengthen the two subproblems in  $Y, R$ . In addition, we periodically find lower and upper bound estimates in order to stop the algorithm early, i.e., with low accuracy.

Our numerical experiments show that our approach for solving **MC** improves on the existing approach in [19]. And furthermore it is also competitive with the approach of [22] in the context of the vertex separator problem.

## References

1. MOSEK ApS. *The MOSEK optimization toolbox for MATLAB manual. Version 8.1.*, (2017). <http://docs.mosek.com/8.0/toolbox/index.html>
2. Balas, E., Ceria, S., Cornuejols, G.: A lift-and-project cutting plane algorithm for mixed 0–1 programs. *Math. Programm.* **58**, 295–324 (1993)
3. Boyd, S., Parikh, N., Chu, E., Peleato, B., Eckstein, J.: Distributed optimization and statistical learning via the alternating direction method of multipliers. *Found. Trends Mach. Learn.* **3**(1), 1–122 (2011)
4. Bücker, H.M., Rostami, M.A.: Interactively exploring the connection between nested dissection orderings for parallel cholesky factorization and vertex separators. In: 2014 IEEE International Parallel Distributed Processing Symposium Workshops, pp. 1122–1129 (2014)
5. Chazelle, Bernard; Kingsford, Carl, Singh, Mona: a semidefinite programming approach to side chain positioning with new rounding strategies. *INFORMS J. Comput.* **16**(4), 380–392 (2004)
6. Chen, Y., Ye, X.: Projection onto a simplex. arXiv preprint. [arXiv:1101.6081](https://arxiv.org/abs/1101.6081), (2011)
7. Cornaz, D., Magnouche, Y., Mahjoub, A.R., Martin, S.: The multi-terminal vertex separator problem: polyhedral analysis and branch-and-cut. *Dis. Appl. Math.* **256**, 11–37 (2019)
8. Didi Biha, M., Meurs, M.-J.: An exact algorithm for solving the vertex separator problem. *J. Global Optim.* **49**(3), 425–434 (2011)
9. George, Alan, Gilbert, John R., Liu, Joseph W.H.: *Graph theory and sparse matrix computation*, vol. 56. Springer, Berlin (2012)
10. Hager, W.W., Hungerford, J.T., Safro, I.: A multilevel bilinear programming algorithm for the vertex separator problem. *Comput. Optim. Appl.* **69**(1), 189–223 (2018)
11. He, B., Liu, H., Wang, Z., Yuan, X.: A strictly contractive Peaceman-Rachford splitting method for convex programming. *SIAM J. Optim.* **24**(3), 1011–1040 (2014)
12. He, Bingsheng; Ma, Feng, Yuan, Xiaoming: convergence study on the symmetric version of ADMM with larger step sizes. *SIAM J. Imag. Sci.* **9**(3), 1467–1501 (2016)
13. Kingsford, Carleton L.; Chazelle, Bernard, Singh, Mona: Solving and analyzing side-chain positioning problems using linear and integer programming. *Bioinformatics* **21**(7), 1028–1039 (2005)
14. Lara, Carlos, Flores, Juan J., Calderon, Felix: On the hyperbox-hyperplane intersection problem. *INFOCOMP J. Comp. Sci.* **8**(4), 21–27 (2009)
15. Lewis, R.H.: Yet another graph partitioning problem is NP-hard. Technical report, (2014)
16. Lovász, L., Schrijver, A.: Cones of matrices and set-functions and 0–1 optimization. *SIAM J. Optim.* **1**(2), 166–190 (1991)
17. Maculan, N., Santiago, C.P., Macambira, E.M., Jardim, M.H.C.: An  $O(n)$  algorithm for projecting a vector on the intersection of a hyperplane and a box in  $\mathbb{R}^n$ . *J. Optim. Theory Appl.* **117**(3), 553–574 (2003)
18. Oliveira, D.E., Wolkowicz, H., Xu, Y.: ADMM for the SDP relaxation of the QAP. *Math. Program. Comput.* **10**(4), 631–658 (2018)
19. Pong, T.K., Sun, H., Wang, N., Wolkowicz, H.: Eigenvalue, quadratic programming, and semidefinite programming relaxations for a cut minimization problem. *Comput. Optim. Appl.* **63**(2), 333–364 (2016)
20. Pothén, A.: Graph partitioning algorithms with applications to scientific computing. In: Keyes, D.E., Sameh, A., Venkatakrishnan, V. (eds.) *Parallel Numerical Algorithms*. ICASE/LaRC Interdisciplinary Series in Science and Engineering, vol. 4, pp. 323–368. Kluwer Academic Press, Dordrecht (1997)
21. Rendl, F., Lisser, A., Piacentini, M.: Bandwidth, vertex separators and eigenvalue optimization. In: Bezdek, K., Deza, A., Ye, Y. (eds.) *Discrete Geometry and Optimization*. Fields Institute Communications, vol. 69, pp. 29–263. Springer (2013)

22. Rendl, F., Sotirov, R.: The min-cut and vertex separator problem. *Comput. Optim. Appl.* **69**(1), 159–187 (2018)
23. Robinson, S.M.: Regularity and stability for convex multivalued functions. *Math. Oper. Res.* **1**, 130–143 (1976)
24. Rockafellar, R.T.: *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks
25. Sherali, H.D., Adams, W.P.: Computational advances using the reformulation-linearization technique (RLT) to solve discrete and continuous nonconvex problems. *Optima* **49**, 1–6 (1996)
26. Simpson, Toby: Sparse cholesky reordering with the graph  $p$ -laplacian. Master's thesis, Università della Svizzera Italiana, (2017)
27. Sun, H.: ADMM for SDP relaxation of GP. Master's thesis, University of Waterloo, (2016)
28. van den Berg, E., Friedlander, M.P.: Probing the Pareto frontier for basis pursuit solutions. *SIAM J. Sci. Comput.* **31**(2), 890–912 (2008)
29. Wolkowicz, H., Zhao, Q.: Semidefinite programming relaxations for the graph partitioning problem. *Discrete Appl. Math.*, 96/97:461–479, 1999. Selected for the special Editors' Choice, Edition (1999)
30. Zhao, Q.: Semidefinite Programming for Assignment and Partitioning Problems. PhD thesis, University of Waterloo, (1996)
31. Zhao, Q., Karisch, S.E., Rendl, F., Wolkowicz, H.: Semidefinite programming relaxations for the quadratic assignment problem. *J. Comb. Optim.* **2**, 71–109 (1998)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliation

Xinxin Li<sup>1</sup> · Ting Kei Pong<sup>2</sup> · Hao Sun<sup>3</sup> · Henry Wolkowicz<sup>3</sup> 

✉ Henry Wolkowicz  
hwolkowicz@uwaterloo.ca

Xinxin Li  
xinxinli@jlu.edu.cn

Ting Kei Pong  
tk.pong@polyu.edu.hk

Hao Sun  
h34sun@uwaterloo.ca

<sup>1</sup> School of Mathematics, Jilin University, Changchun, China

<sup>2</sup> Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong, The People's Republic of China

<sup>3</sup> Department of Combinatorics & Optimization, University of Waterloo, Waterloo, Canada