

3.1 Introduction to Determinants

Notation: A_{ij} is the matrix obtained from matrix A by deleting the i th row and j th column of A .

EXAMPLE:

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix} \quad A_{23} = \begin{bmatrix} 1 & 2 & 4 \\ 9 & 10 & 12 \\ 13 & 14 & 16 \end{bmatrix}$$

Recall that $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$ and we let $\det[a] = a$.

For $n \geq 2$, the **determinant** of an $n \times n$ matrix $A = [a_{ij}]$ is given by

$$\begin{aligned} \det A &= a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{1+n} a_{1n} \det A_{1n} \\ &= \sum_{j=1}^n (-1)^{1+j} a_{1j} \det A_{1j} \end{aligned}$$

EXAMPLE: Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$

Solution

$$\begin{aligned} \det A &= 1 \det \begin{bmatrix} -1 & 2 \\ 0 & 1 \end{bmatrix} - 2 \det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} + 0 \det \begin{bmatrix} 3 & -1 \\ 2 & 0 \end{bmatrix} \\ &= \underline{1(-1) - 2(-1) + 0(2)} = \underline{1} \end{aligned}$$

Common notation: $\det \begin{bmatrix} 3 & 2 \\ 2 & 1 \end{bmatrix} = \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix}$.

So

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1 \begin{vmatrix} -1 & 2 \\ 0 & 1 \end{vmatrix} - 2 \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix}$$

The **(i, j)-cofactor** of A is the number C_{ij} where $C_{ij} = (-1)^{i+j} \det A_{ij}$.

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 1C_{11} + 2C_{12} + 0C_{13}$$

(cofactor expansion across row 1)

THEOREM 1 The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column:

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in} \quad (\text{expansion across row } i)$$

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj} \quad (\text{expansion down column } j)$$

Use a matrix of signs to determine $(-1)^{i+j}$

$$\begin{bmatrix} + & - & + & \cdots \\ - & + & - & \cdots \\ + & - & + & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

EXAMPLE: Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$

using cofactor expansion down column 3.

Solution

$$\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} = 0 \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 2 & 0 \end{vmatrix} + 1 \begin{vmatrix} 1 & 2 \\ 3 & -1 \end{vmatrix} = 1.$$

EXAMPLE: Compute the determinant of $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{bmatrix}$

Solution

$$\begin{vmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 3 & 5 \end{vmatrix} \quad (\text{cofactor expansion along column 1})$$

$$= 1 \begin{vmatrix} 2 & 1 & 5 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 0 & 2 & 1 \\ 0 & 3 & 5 \end{vmatrix} + 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 2 & 3 & 4 \\ 2 & 1 & 5 \\ 0 & 2 & 1 \end{vmatrix}$$

$$= 1 \cdot 2 \begin{vmatrix} 2 & 1 \\ 3 & 5 \end{vmatrix} = 14$$

Method of cofactor expansion is not practical for large matrices - see Numerical Note on page 190.

Triangular Matrices:

$$\begin{bmatrix} * & * & \cdots & * & * \\ 0 & * & \cdots & * & * \\ 0 & 0 & \ddots & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & * \end{bmatrix}$$

(upper triangular)

$$\begin{bmatrix} * & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 \\ * & * & \ddots & 0 & 0 \\ * & * & \cdots & * & 0 \\ * & * & \cdots & * & * \end{bmatrix}$$

(lower triangular)

THEOREM 2: If A is a triangular matrix, then $\det A$ is the product of the main diagonal entries of A .

EXAMPLE:

$$\begin{vmatrix} 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & -3 & 5 \\ 0 & 0 & 0 & 4 \end{vmatrix} = \frac{2(1)(-3)(4)}{1} = -24$$