

NETWORK DESIGN VIA ITERATIVE ROUNDING  
OF SETPAIR RELAXATIONS

JOSEPH CHERIYAN\*, SANTOSH VEMPALA†, ADRIAN VETTA

Received August 16, 2001

Revised October 2, 2001

A typical problem in network design is to find a minimum-cost sub-network  $H$  of a given network  $G$  such that  $H$  satisfies some prespecified connectivity requirements. Our focus is on approximation algorithms for designing networks that satisfy *vertex connectivity* requirements. Our main tool is a linear programming relaxation of the following *setpair formulation* due to Frank and Jordan: a *setpair* consists of two subsets of vertices (of the given network  $G$ ); each setpair has an integer requirement, and the goal is to find a minimum-cost subset of the edges of  $G$  such that each setpair is covered by at least as many edges as its requirement. We introduce the notion of *skew bisupermodular* functions and use it to prove that the basic solutions of the linear program are characterized by “non-crossing families” of setpairs. This allows us to apply Jain’s *iterative rounding method* to find approximately optimal integer solutions. We give two applications. (1) In the *k-vertex connectivity problem* we are given a (directed or undirected) graph  $G = (V, E)$  with nonnegative edge costs, and the task is to find a minimum-cost spanning subgraph  $H$  such that  $H$  is  $k$ -vertex connected. Let  $n = |V|$ , and let  $\epsilon < 1$  be a positive number such that  $k \leq (1 - \epsilon)n$ . We give an  $O(\sqrt{n/\epsilon})$ -approximation algorithm for both problems (directed or undirected), improving on the previous best approximation guarantees for  $k$  in the range  $\Omega(\sqrt{n}) < k < n - \Omega(1)$ . (2) We give a 2-approximation algorithm for the *element connectivity* problem, matching the previous best approximation guarantee due to Fleischer, Jain and Williamson.

---

*Mathematics Subject Classification (2000):* 68W25, 90C35; 05C40, 68R10, 90C27, 90B10

\* Supported in part by NSERC research grant OGP0138432.

† Supported in part by NSF Career Award CCR-9875024.

## 1. Introduction

A typical problem in network design is to find a minimum-cost sub-network  $H$  of a given network  $G$  such that  $H$  satisfies some prespecified connectivity requirements; examples are the *minimum spanning tree* (MST) problem and the *traveling salesman problem* (TSP). By a *network* we mean a graph (either undirected or directed) together with non-negative costs for the edges. Problems in network design have a central position in combinatorial optimization and theoretical computer science, and moreover, they arise in many practical settings (e.g., telephone networks). Many of the problems in network design are NP-hard. Over the past decade, there has been significant research on approximation algorithms for network design, and there have been some notable successes in the design of networks that satisfy various types of “edge connectivity” requirements, e.g., Goemans and Williamson [9], Jain [11], etc. Fewer results are known on the design of networks subject to “vertex connectivity” requirements. There are several recent results for restricted edge costs, such as uniform costs or metric costs, but in this paper we discuss only the general case of non-negative costs. Our focus is on approximation algorithms for designing networks that satisfy vertex connectivity requirements. Our results are based on the iterative rounding method (due to Jain [11]) and on an integer programming formulation called the *setpair formulation* (due to Frank and Jordan [5]).

Let  $G = (V, E)$  be the graph of the given network, and let  $n$  denote the number of vertices. Most of the previous research in this area is based on an integer programming formulation called the *cut covering* model. A non-negative, integer requirement  $f(S, V \setminus S)$  is assigned to each vertex partition  $(S, V \setminus S)$ , and the goal is to find a minimum-cost subgraph  $H$  such that each cut  $(S, V \setminus S)$  has at least  $f(S, V \setminus S)$  edges of  $H$ . (For example, in the MST problem,  $f(S, V \setminus S) = 1$  for each nonempty vertex set  $S$ ,  $S \neq V$ .) Although this model captures many “edge connectivity” problems in network design, it seems less suited for “vertex connectivity” problems.

The *iterative rounding* method works as follows. Formulate the problem as an integer program, and then solve the LP (linear programming) relaxation to find a basic (extreme point) optimum solution  $\mathbf{x}$ . Pick an edge  $e^*$  of highest value (i.e.,  $x_{e^*} \geq x_e, \forall e \in E$ ) and add it to the solution subgraph  $H$  (initially,  $E(H)$  is empty). Then update the LP and the integer program, since the variable  $x_{e^*}$  is implicitly fixed at value 1. In detail, decrease by 1 the r.h.s. of every constraint where the variable  $x_{e^*}$  occurs, and then remove this variable from the LP. The resulting LP is the same as the LP for the “reduced” problem where the edge  $e^*$  is pre-selected for  $H$ . Under appropriate conditions on the requirement function  $f$ , the problem turns out to be “self

reducible,” i.e., the essential properties of the original problem are preserved in the reduced problem. Iteratively solve the reduced problem. Jain [11] applied this method to the cut covering model, and proved that it achieves an approximation guarantee of 2 provided that the requirement function  $f$  is weakly supermodular. (Such requirement functions capture several interesting problems, e.g., the Steiner network problem.) His proof is based on a key property of the LP: every non-zero basic solution has an edge of value at least  $\frac{1}{2}$ . This result is based on an extension of a classic result that, under appropriate conditions on the requirement function  $f$ , every basic solution of the LP is characterized by a laminar family of “tight sets.”

The setpair formulation [5] is a generalization of the cut covering model. A setpair  $(W_t, W_h)$  consists of two disjoint vertex sets  $W_t$  and  $W_h$ , i.e.,  $W_t \subseteq V$ ,  $W_h \subseteq V$ ,  $W_t \cap W_h = \emptyset$ . Each setpair  $(W_t, W_h)$  is assigned a non-negative, integer requirement  $f(W_t, W_h)$ . The goal is to find a minimum-cost subgraph  $H$  that satisfies the requirement of every setpair, i.e., for each setpair  $(W_t, W_h)$ ,  $H$  should have at least  $f(W_t, W_h)$  edges that have one end-vertex in  $W_t$  and the other end-vertex in  $W_h$ . Note that the cut covering model is a special case of the setpair formulation in which every setpair  $(W_t, W_h)$  with a positive requirement has  $W_t = V \setminus W_h$  (i.e., a pair of complementary sets).

We introduce a special class of requirement functions  $f$ , namely, *skew bisupermodular* functions (see Section 2). These functions are a common generalization of weakly supermodular (symmetric) functions ([3], [8]) and crossing bisupermodular functions [5]. Skew bisupermodular requirement functions are useful because the basic solutions of the LP relaxation are characterized by “non-crossing families” of setpairs (see Section 3 and Theorem 3.3). Note the correspondence with the cut covering model, where the basic solutions of the LP relaxation may be characterized by a laminar family of sets. Based on this, we obtain lower bounds on the maximum value of an edge for the basic solutions of the LP relaxation. Also, this LP relaxation has the self-reducibility property needed for the iterative rounding method. Hence, these structural results immediately give approximation algorithms.

We give applications to two specific problems. In the *k-vertex connectivity problem* we are given a graph  $G = (V, E)$  (either directed or undirected) with nonnegative edge costs  $c: E \rightarrow \mathbb{R}_+$ , and the task is to find a minimum-cost spanning subgraph  $H$  such that  $H$  is *k-vertex connected*. A graph is called *k-vertex connected* if it has at least  $k+1$  vertices, and the deletion of any set of  $k-1$  vertices leaves a connected graph. We give an approximation algorithm for the undirected problem, and our method extends to the directed problem. Our approximation guarantee is  $O(\sqrt{n/\epsilon})$  where  $\epsilon$  is a positive real number

such that  $k \leq (1 - \epsilon)n$ . For both the directed and the undirected problems, our results improve on the previous best approximation guarantees provided that  $k$  is in the range  $\Omega(\sqrt{n}) < k < n - \Omega(1)$ . The previous best approximation guarantee for the directed problem was  $\Omega(k)$  [6], and the previous best approximation guarantee for the undirected problem was  $O(\log k)$  for  $k \leq \sqrt{\frac{n}{6}}$  [2], and  $\Omega(k)$  for  $k > \sqrt{\frac{n}{6}}$  [14]. (An  $O(\log k)$  approximation guarantee for the undirected problem was claimed earlier in [15], but subsequently an error has been found and that claim has been withdrawn.) Very recently (and two years after this paper was completed), improved approximation guarantees for the  $k$ -vertex connectivity problem have been reported in [13].

In the *element connectivity* problem [12], we are given an undirected graph  $G = (V, E)$  with nonnegative edge costs  $c: E \rightarrow \mathbb{R}_+$ ; moreover, there is a set of *terminal* vertices  $T \subseteq V$  and between each pair of distinct terminals  $i$  and  $j$  a connectivity requirement  $r_{ij}$  is specified (each  $r_{ij}$  is a non-negative integer, and  $r_{ij} = r_{ji}$ ). Terminals are reliable and do not fail. Edges and non-terminal vertices are unreliable and are subject to failure. The edges and non-terminals are called *elements*. The problem is to find a minimum-cost subgraph  $H$  that contains  $r_{ij}$  element disjoint paths between each pair of terminals  $i$  and  $j$ . Alternatively, for every pair of terminals  $i$  and  $j$ ,  $H$  should contain a path between  $i$  and  $j$  despite the failure of up to  $r_{ij} - 1$  elements. The problem is NP-hard. Recently, Fleischer, Jain, and Williamson [7] gave a 2-approximation algorithm for this problem, improving on the previous best approximation guarantee of  $2 \ln(\max\{r_{ij}\})$  due to Jain et al [12]. We give a different proof of the same 2-approximation guarantee. The element connectivity problem is a generalization of the Steiner network problem, see Goemans et al [8] and Jain [11]. Even for special cases of this problem, the best approximation guarantee known is 2, and moreover, the LP relaxation has an integrality ratio of at least  $2 - \frac{2}{n}$ .

Let us briefly comment on the relation of this work to [7]. We use the same linear programming relaxation for the element connectivity problem as Fleischer et al, but the two proofs are different. In particular, the definitions of “noncrossing setpairs” are different. (In our setting, noncrossing setpairs may be tail disjoint, or head disjoint, or comparable (this follows [1, 5]), whereas [7] has a more restricted notion of noncrossing setpairs.) We have included a proof of (a generalization of) the 2-approximation guarantee for element connectivity for the sake of completeness.

For many of the specific problems in network design that are captured by the setpairs formulation, the linear programming relaxation can be written as a compact linear program via a “flow formulation.” Moreover, an appropriate optimal solution can be computed in strongly polynomial time via

Tardos' algorithm [16]. The computed optimal solution may not be basic, but it will have an edge of sufficiently large value. This applies to both the  $k$ -vertex connectivity problem and the element connectivity problem, and is similar to the method used by Jain in [11, Section 9].

Summarizing, our goal is to present general results on the iterative rounding of setpair relaxations, to model diverse problems in network design, and to derive good approximation guarantees for specific problems. The rest of this paper is organized as follows. Section 2 has notation and definitions pertaining to the setpair formulation. Section 3 contains our main result on skew bisupermodular functions, a combinatorial characterization of the basic solutions of the LP. Section 4 gives the approximation algorithm for the  $k$ -vertex connectivity problem. The concluding section, Section 5 describes a setpair formulation for the element connectivity problem, and gives our proof of the 2-approximation guarantee.

## 2. The Setpair Formulation and Skew Bisupermodular Functions

One of the new contributions of this section is the notion of skew bisupermodular functions (and the related notions of reverse setpairs and overlapping setpairs). Most of the other material is based on [5].

A *setpair*  $W = (W_t, W_h)$  is an ordered pair of disjoint vertex sets; either  $W_t$  or  $W_h$  may be empty. We say that  $W_t$  is the *tail* of  $W$ , and  $W_h$  is the *head* of  $W$ . Let  $\mathcal{S}$  denote the set of all setpairs. Denote by  $\delta(W) = \delta(W_t, W_h)$  the set of edges with one end-vertex in  $W_t$  and the other in  $W_h$ ; note that a *cut*  $\delta(W_t, V \setminus W_t)$  is the special case of  $\delta(W)$  where the head and tail of  $W$  form a complementary pair of vertex sets.

The setpair formulation (discussed in Section 1) may be written as the following integer program (IP).

$$\begin{array}{ll}
 & \text{minimize} & \sum_{e \in E} c_e x_e \\
 \text{(IP)} & \text{subject to} & \sum_{e \in \delta(W)} x_e \geq f(W), & \forall W \in \mathcal{S} \\
 & & x_e \in \{0, 1\}, & \forall e \in E
 \end{array}$$

Here,  $f$  is some non-negative integer function on the setpairs, and  $c: E \rightarrow \mathbb{R}_+$  gives the edge costs. Let (LP) denote the linear programming relaxation obtained from (IP) by replacing the constraints  $x_e \in \{0, 1\}$  by  $0 \leq x_e \leq 1$  for all edges  $e \in E$ .

### 2.1. Crossing Setpairs.

Given two setpairs  $W = (W_t, W_h)$  and  $Y = (Y_t, Y_h)$ , let  $W \otimes Y$  denote the setpair  $(W_t \cup Y_t, W_h \cap Y_h)$ , and let  $W \oplus Y$  denote the setpair  $(W_t \cap Y_t, W_h \cup Y_h)$ .

Two setpairs  $W$  and  $Y$  are said to be *comparable* if

- (i)  $W_t \supseteq Y_t$  and  $W_h \subseteq Y_h$ , or (ii)  $W_t \subseteq Y_t$  and  $W_h \supseteq Y_h$ .

The former case is denoted by  $W \preceq Y$  and the latter case by  $W \succeq Y$ . Note that  $W \preceq W \oplus Y$  and  $W \succeq W \otimes Y$ .

Two setpairs  $W$  and  $Y$  are said to be *non-crossing* if

- (i) they are comparable, or (ii) their tails are disjoint, or (iii) their heads are disjoint.

Otherwise,  $W$  and  $Y$  are said to *cross*. Note that  $W$  and  $Y$  cross if and only if their tails intersect, their heads intersect and they are not comparable.

For a setpair  $W = (W_t, W_h)$ , let  $\overline{W} = (W_h, W_t)$  be the *reverse setpair*. Observe that  $\delta(W) = \delta(\overline{W})$  for an undirected graph. Also, note that  $\overline{W \otimes Y} = \overline{W} \oplus \overline{Y}$ .

Two setpairs  $W$  and  $Y$  are said to be *overlapping* if

- (i)  $W$  and  $Y$  cross, or (ii)  $\overline{W}$  and  $Y$  cross (which is the same as  $W$  and  $\overline{Y}$  cross).

Otherwise, the setpairs are called *non-overlapping*. A family of setpairs  $\mathcal{L} \subseteq \mathcal{S}$  is called non-crossing (non-overlapping) if no two setpairs in  $\mathcal{L}$  cross (overlap).

### 2.2. Bisubmodular Functions.

A real-valued function  $f$  on  $\mathcal{S}$  is called *bisubmodular* if for any two setpairs  $W$  and  $Y$  we have

$$f(W) + f(Y) \geq f(W \otimes Y) + f(W \oplus Y).$$

A non-negative, integer-valued function  $f$  on  $\mathcal{S}$  is called *crossing bisupermodular* if for any two crossing setpairs  $W$  and  $Y$  with  $f(W) > 0$  and  $f(Y) > 0$ , we have

$$f(W) + f(Y) \leq f(W \otimes Y) + f(W \oplus Y).$$

Let  $\chi_W$  denote the edge incidence vector of  $\delta(W)$ . For any two setpairs  $W$  and  $Y$ , note that if an edge is present in  $\delta(W \oplus Y)$  or  $\delta(W \otimes Y)$ , then it is present in  $\delta(W)$  or  $\delta(Y)$ , and if an edge is present in both  $\delta(W \oplus Y)$  and  $\delta(W \otimes Y)$ , then it is present in both  $\delta(W)$  and  $\delta(Y)$ . Hence, we have  $\chi_{W \otimes Y} + \chi_{W \oplus Y} \leq \chi_W + \chi_Y$ . Consequently, for any non-negative vector  $\mathbf{x}: E \rightarrow \mathbb{R}_+$  on the edges, the corresponding function on setpairs,  $x(\delta(W)) = \sum_{e \in \delta(W)} x_e$ , is

bisubmodular. (For any vector  $\mathbf{x}$  on a groundset  $U$  and a subset  $Q$  of  $U$ ,  $x(Q)$  denotes  $\sum_{i \in Q} x_i$ .)

### 2.3. Skew Bisupermodular Functions.

A non-negative, integer-valued function  $f$  on  $\mathcal{S}$  is called *skew bisupermodular* if for any two overlapping setpairs  $W$  and  $Y$  with  $f(W) > 0$  and  $f(Y) > 0$ , we have

$$(i) \quad f(W) + f(Y) \leq f(W \otimes Y) + f(W \oplus Y), \quad \text{or}$$

$$(ii) \quad f(W) + f(Y) \leq f(\overline{W} \otimes Y) + f(\overline{W} \oplus Y).$$

This is motivated by problems on undirected graphs. For these problems, it turns out that the relevant functions  $f$  are symmetric, i.e.,  $f(W) = f(\overline{W})$ ,  $\forall W \in \mathcal{S}$ .

Note that every crossing bisupermodular function is skew bisupermodular. Also, every weakly supermodular symmetric function is skew bisupermodular (see the proof of Lemma 5.2). Thus the skew bisupermodular property is a generalization of these earlier notions.

### 3. Characterizing a Basic Solution via a Non-Overlapping Family

The main result of this section, Theorem 3.3, characterizes a basic solution of (LP) via a non-overlapping family of setpairs, assuming that the given graph  $G$  is undirected and the requirement function  $f$  is symmetric and skew bisupermodular. This result is an extension of classic results to our setting. Our proof is similar to earlier proofs in [11, Lemma 4.2] and [1, Theorem 2.1]; however, when we add a new setpair  $W$  to our non-overlapping family, we have to ensure that neither  $W$  nor  $\overline{W}$  crosses a setpair in the family. The results in this section may not apply to directed graphs.

Given a feasible solution  $\mathbf{x}$  to (LP), a setpair  $W$  is called *tight* if  $x(\delta(W)) = f(W)$ .

**Lemma 3.1.** *Let the requirement function  $f$  of (LP) be symmetric and skew bisupermodular, and let  $\mathbf{x}$  be a feasible solution to (LP) such that  $x_e > 0$  for all edges  $e \in E$ . Suppose that the setpairs  $W$  and  $Y$  have  $f(W) > 0$ ,  $f(Y) > 0$ , and moreover,  $W$  and  $Y$  overlap, and are tight (also, note that  $\overline{W}$  is tight, it overlaps  $Y$ , and  $f(\overline{W}) > 0$ ). Then one of the following holds:*

- (i) *The setpairs  $W \otimes Y$  and  $W \oplus Y$  are tight, and  $\chi_W + \chi_Y = \chi_{W \otimes Y} + \chi_{W \oplus Y}$ .*
- (ii) *The setpairs  $\overline{W} \otimes Y$  and  $\overline{W} \oplus Y$  are tight, and  $\chi_W + \chi_Y = \chi_{\overline{W} \otimes Y} + \chi_{\overline{W} \oplus Y}$ .*

**Proof.** Assume that  $f(W) + f(Y) \leq f(W \otimes Y) + f(W \oplus Y)$ . The other case is similar. Then:

$$\begin{aligned} f(W \otimes Y) + f(W \oplus Y) &\leq x(\delta(W \otimes Y)) + x(\delta(W \oplus Y)) \\ &\leq x(\delta(W)) + x(\delta(Y)) \\ &= f(W) + f(Y) \\ &\leq f(W \otimes Y) + f(W \oplus Y) \end{aligned}$$

Hence, all the inequalities hold as equations, therefore,  $W \otimes Y$  and  $W \oplus Y$  are both tight. This proves the first part of (i). The second part of (i) follows from the following three facts:  $\chi_W + \chi_Y \geq \chi_{W \otimes Y} + \chi_{W \oplus Y}$ ;  $x_e > 0$  for all edges  $e$ ;  $x(\delta(W \otimes Y)) + x(\delta(W \oplus Y)) = x(\delta(W)) + x(\delta(Y))$ . ■

The next result is in [1, Lemma 2.3] and is used to prove the main result of this section.

**Lemma 3.2.** *Let  $W$ ,  $Y$  and  $Z$  be setpairs. If  $Z$  crosses  $W \otimes Y$  (or  $W \oplus Y$ ) then either  $Z$  crosses  $W$  or  $Z$  crosses  $Y$ .* ■

**Theorem 3.3.** *Let the requirement function  $f$  of (LP) be symmetric and skew bisupermodular, and let  $\mathbf{x}$  be a basic solution of (LP) such that  $0 < x_e < 1$  for all edges  $e \in E$ . Then there exists a non-overlapping family  $\mathcal{L}$  of tight setpairs such that:*

- (i) Every setpair  $W \in \mathcal{L}$  has  $f(W) \geq 1$ .
- (ii)  $|\mathcal{L}| = |E|$ .
- (iii) The vectors  $\chi_W$ ,  $W \in \mathcal{L}$ , are linearly independent.
- (iv)  $\mathbf{x}$  is the unique solution to  $\{x(\delta(W)) = f(W), \forall W \in \mathcal{L}\}$ .

**Proof.** Since  $\mathbf{x}$  is a basic solution of (LP), and none of the constraints  $0 \leq x_e \leq 1$  ( $e \in E$ ) holds with equality, there exist  $|E|$  tight setpair constraints  $x(\delta(W)) = f(W) > 0$  such that the edge incidence vectors  $\chi_W$  corresponding to these constraints are linearly independent. Let  $\mathcal{L}$  be a maximal, non-overlapping family of tight setpairs and let  $\text{span}(\mathcal{L})$  denote the vector space spanned by the vectors  $\chi_W$ , where  $W \in \mathcal{L}$ . The theorem holds because  $\text{span}(\mathcal{L})$  equals  $\text{span}(\mathcal{F})$ , where  $\mathcal{F}$  is the family of all tight setpairs. We prove this by contradiction. Suppose that there is a tight setpair  $Y$  such that  $\chi_Y \notin \text{span}(\mathcal{L})$ . Take such a  $Y$  that overlaps the minimum number of setpairs in  $\mathcal{L}$  (this is a key step). Choose any setpair  $W \in \mathcal{L}$  that overlaps  $Y$ ; there exists such a  $W$  by the maximality of  $\mathcal{L}$ .

Either part (i) or part (ii) in the statement of Lemma 3.1 holds for  $W$  and  $Y$ . Assume that part (ii) holds. The other case is similar. Thus, the setpairs  $\overline{W \otimes Y}$  and  $\overline{W \oplus Y}$  are tight. In addition,  $\chi_W + \chi_Y = \chi_{\overline{W \otimes Y}} + \chi_{\overline{W \oplus Y}}$ . It



follows that either  $\chi_{\overline{W} \otimes Y} \notin \text{span}(\mathcal{L})$  or  $\chi_{\overline{W} \oplus Y} \notin \text{span}(\mathcal{L})$ . Consider the former case. Again, the other case is similar. Let  $Q = \overline{W} \otimes Y$  and take any setpair  $Z \in \mathcal{L}$ . Suppose that  $Z$  overlaps  $Q$ . Then either  $Z$  crosses  $Q$  or  $Z$  crosses  $\overline{Q} = \overline{\overline{W} \otimes Y} = \overline{W} \oplus \overline{Y}$ . Suppose that  $Z$  crosses  $Q$ . We apply Lemma 3.2 to  $Z$  and  $Q$ , noting that  $Z$  does not cross  $\overline{W}$ , to see that  $Z$  crosses  $Y$ . Similarly, if  $Z$  crosses  $\overline{Q}$ , then by Lemma 3.2,  $Z$  crosses  $\overline{Y}$ . Hence,  $Z$  overlaps  $Y$ . So every setpair in  $\mathcal{L}$  that overlaps  $Q$  also overlaps  $Y$ .

It follows that  $Q$  overlaps fewer setpairs in  $\mathcal{L}$  than  $Y$ . To see this, note that (1)  $W$  overlaps  $Y$ , by our choice of  $W$ , and (2)  $W$  does not overlap  $Q$ , because  $Q \preceq \overline{W}$  (so  $Q$  does not cross  $\overline{W}$ ) and  $W$  and  $Q$  are head disjoint (so  $W$  does not cross  $Q$ ). Thus we get a contradiction to the choice of  $Y$ , since  $Q$  is a tight setpair that overlaps fewer setpairs in  $\mathcal{L}$  than  $Y$ , and  $\chi_Q \notin \text{span}(\mathcal{L})$ . ■

**Remarks.** In the theorem, if we replace a setpair  $W \in \mathcal{L}$  by its reverse  $\overline{W}$ , then this preserves the properties (i)–(iv).

#### 4. An Approximation Algorithm for the $k$ -Vertex Connectivity Problem

The main result of this section is as follows.

**Theorem 4.1.** *Let  $k$  and  $n$  be positive integers, and let  $\epsilon < 1$  be a positive number such that  $k$  is at most  $(1-\epsilon)n$ . There is a polynomial-time algorithm that, given an  $n$ -vertex (directed or undirected) graph, finds a solution to the  $k$ -vertex connectivity problem of cost at most  $O(\sqrt{n/\epsilon})$  times the optimal cost.* ■

The  $k$ -vertex connectivity problem is modeled by the following integer program.

$$\begin{aligned}
 \text{(IP-VC)} \quad & \text{minimize} && \sum_{e \in E} c_e x_e \\
 & \text{subject to} && \sum_{e \in \delta(W)} x_e \geq f(W) && \forall W \in \mathcal{S} \\
 & && x_e \in \{0, 1\} && \forall e \in E
 \end{aligned}$$

Here,  $f$  is the following non-negative integer function defined on the setpairs.

$$f(W) = \begin{cases} \max \{0, k - |V \setminus (W_h \cup W_t)|\}, & \text{if } W_t \neq \emptyset \text{ and } W_h \neq \emptyset; \\ 0, & \text{otherwise.} \end{cases}$$

We use (LP-VC) to denote the linear programming relaxation.

The  $k$ -vertex connectivity requirement function  $f$  is crossing bisupermodular [5], hence,  $f$  is also skew bisupermodular. This can be seen as follows. For each setpair  $W$ , let  $p(W)$  denote  $|V \setminus (W_t \cup W_h)|$ . For any two setpairs  $W = (W_t, W_h)$  and  $Y = (Y_t, Y_h)$ , we have  $p(W) + p(Y) = p(W \otimes Y) + p(W \oplus Y)$ , because

$$|(W_t \cup Y_t) \cup (W_h \cap Y_h)| + |(W_t \cap Y_t) \cup (W_h \cup Y_h)| = |W_t| + |W_h| + |Y_t| + |Y_h|.$$

Hence, the above function  $f(\cdot)$  is crossing bisupermodular (if either  $W \otimes Y$  or  $W \oplus Y$  has head or tail empty, then note that  $W, Y$  do not cross). The results in [5] imply that a basic optimal solution to (LP-VC) (if it exists) can be found in polynomial time, see [5], [10, Theorem 6.4.9]. Moreover, the requirement function  $f$  and (LP-VC) have the following “self-reducibility” property: for any edge  $e$  the “reduced” requirement function  $f'$  is also crossing bisupermodular (and skew bisupermodular), where  $f'$  is given by  $f'(W) = f(W) - 1$  if the setpair  $W$  has  $e \in \delta(W)$  and  $f'(W) = f(W)$  otherwise (since the zero-one incidence function of  $e$  on  $\mathcal{S}$  is bisubmodular).

Theorem 4.1 follows directly from the next result.

**Theorem 4.2.** *Let  $\epsilon < 1$  be a positive number, and suppose that  $k \leq (1-\epsilon)n$ . Then any nonzero basic solution of (LP-VC) has an edge of value  $\Omega(\sqrt{\frac{\epsilon}{n}})$ .*

We give the proof for undirected graphs, using results on skew bisupermodular functions. After that, we sketch the proof for directed graphs via crossing bisupermodular functions. The next result is from [1, Theorem 1.1].

**Theorem 4.3.** *Suppose that the requirement function  $f$  for the linear program (LP-VC) is crossing (or, skew) bisupermodular. Let  $\mathbf{x}$  be a nonzero basic solution of (LP-VC), and let  $\mathcal{L}$  be a non-crossing family of setpairs characterizing  $\mathbf{x}$ . Then there exists an edge  $e$  with  $x_e \geq \frac{1}{\Omega(\sqrt{|\mathcal{L}|})}$ . ■*

Focus on a nonzero basic solution  $\mathbf{x}$  of (LP-VC). By Theorem 3.3,  $\mathbf{x}$  corresponds to a noncrossing family of setpairs  $\mathcal{L}$  such that each setpair has a nonempty head and a nonempty tail. We may assume that each setpair  $W \in \mathcal{L}$  has  $|W_h| \geq |W_t|$ , else we may replace  $W$  by  $\bar{W}$  and this preserves all the properties of  $\mathcal{L}$  in Theorem 3.3. Let  $\mathcal{P}$  denote the poset (and the Hasse diagram) representing  $\mathcal{L}$ , and recall that  $\mathcal{P}$  has a node  $W$  for each setpair  $W$  in  $\mathcal{L}$  and it has an arc  $(W, Y)$ , where  $Y \in \mathcal{L}$ , if  $W \preceq Y$  and there is no other  $Z \in \mathcal{L}$  such that  $W \preceq Z \preceq Y$  (we omit arcs implied by transitivity from  $\mathcal{P}$ ); we call  $W$  a *child* of  $Y$ , and call  $Y$  a *parent* of  $W$ . (Note that the terms “node” and “arc” refer to the poset  $\mathcal{P}$ , and the terms “vertex” and

“edge” refer to the input graph or digraph  $G$ .) A *diamond* of a poset is a set of four nodes  $W, X, Y, Z$  such that  $X$  and  $Y$  are incomparable,  $W$  is a common descendant of  $X, Y$ , and  $Z$  is a common ancestor of  $X, Y$ . A poset is called *diamond-free* if it contains no diamond. The poset  $\mathcal{P}$  representing  $\mathcal{L}$  is diamond-free, see [1, Lemma 3.1].

Let  $m$  denote  $|\mathcal{L}|$ . If  $m < 4n/\epsilon$ , then by the above theorem, there is an edge  $e$  with  $x_e \geq \sqrt{\frac{\epsilon}{\Theta(n)}}$ . Hence, in this case, Theorem 4.2 holds. The rest of the proof focuses on the other case ( $m \geq 4n/\epsilon$ ) and shows that there exists an edge  $e$  with either  $x_e \geq \frac{1}{\Theta(1)}$  or  $x_e \geq \Omega(\epsilon) \geq \Omega(\sqrt{\frac{\epsilon}{n}})$ . (Note that  $k \leq n-1$ , hence  $\epsilon \geq \frac{1}{n}$ , and so  $\epsilon \geq \sqrt{\frac{\epsilon}{n}}$ .)

**Lemma 4.4.** *Suppose that  $k \leq (1-\epsilon)n$ . Then there are at most  $\frac{2}{\epsilon}$  pairwise incomparable setpairs in  $\mathcal{L}$  such that all their tails contain a common vertex.*

**Proof.** Let the setpairs  $Y^1, Y^2, \dots, Y^q \in \mathcal{L}$  be pairwise incomparable, and let  $v$  be a vertex that occurs in the tail of each of these setpairs. Since the  $Y^i$  are incomparable and their tails intersect, their heads are disjoint. Each head  $Y_h^i$  has cardinality at least  $\frac{1}{2}(n-k) \geq \frac{1}{2}\epsilon n$ ; this follows from the properties of  $\mathcal{L}$ , and the fact that each requirement  $f(Y^i)$  is at least one. Hence, there are at most  $\frac{2}{\epsilon}$  such setpairs.  $\blacksquare$

We partition the setpairs in  $\mathcal{L}$  into several sets.

- $W$  is a type I setpair if it has no parents in  $\mathcal{P}$ .
- $W$  is a type II setpair if it has at least two parents in  $\mathcal{P}$ .
- $W$  is a type III setpair if it has exactly one parent  $Y$  in  $\mathcal{P}$ , and  $W_t = Y_t$ .
- $W$  is a type IV setpair if it has exactly one parent  $Y$  in  $\mathcal{P}$ , and  $W_t \neq Y_t$ .

**Lemma 4.5.** *Suppose that  $k \leq (1-\epsilon)n$ . (a) Then there are at most  $\frac{2n}{\epsilon}$  setpairs of type IV. (b) Moreover, each setpair (node) in  $\mathcal{P}$  has at most  $\frac{2}{\epsilon}$  children.*

**Proof.** We prove (a) first, then (b). To each type IV setpair  $W$  with parent  $Y$  we assign a vertex  $v$  in  $W_t \setminus Y_t$ . We claim that a vertex can be assigned to at most  $\frac{2}{\epsilon}$  setpairs of type IV. This will imply the claim. Take a vertex  $v$  that is assigned to the type IV setpairs  $W^1, W^2, \dots, W^q$ . Let their parents be  $Y^1, Y^2, \dots, Y^q$ , respectively. Now  $v \in W_t^1 \cap W_t^2 \cap \dots \cap W_t^q$ , but  $v \notin Y_t^1 \cup Y_t^2 \cup \dots \cup Y_t^q$ . The setpairs  $W^1, W^2, \dots, W^q$  associated with  $v$  must be pairwise incomparable (since  $v$  is in the tail of every  $W^i$ , but for every proper ancestor of a  $W^i$  note that  $v$  is not in the tail). Then by Lemma 4.4 applied to  $W^1, W^2, \dots, W^q$ , there are at most  $\frac{2}{\epsilon}$  such setpairs whose tails contain  $v$ .

For (b), consider all the children  $Y^1, Y^2, \dots, Y^q$  of a node  $W \in \mathcal{P}$ . Then  $Y^1, Y^2, \dots, Y^q$  are pairwise incomparable (otherwise, one of the arcs  $(Y^i, W)$  in  $\mathcal{P}$  is redundant), and moreover, each of the tails  $Y_t^i$ ,  $i=1, \dots, q$ , contains  $W_t$  which is nonempty. Then by Lemma 4.4 applied to  $Y^1, Y^2, \dots, Y^q$ , we have  $q \leq \frac{2}{\epsilon}$ .  $\blacksquare$

We have  $|\mathcal{L}| = m \geq \frac{4n}{\epsilon}$ , hence, by Lemma 4.5,  $\mathcal{L}$  has at least  $\frac{m}{6}$  setpairs of one of the types I, II, or III.

(Type I) Suppose that there are at least  $\frac{m}{6}$  type I setpairs. The type I setpairs are pairwise incomparable. Hence the edge sets  $\delta(W)$ , where  $W$  is a type I setpair, are *quasidisjoint*, i.e., for each edge  $uv$ , there are at most two setpairs  $W, Y$  (among the type I setpairs) such that  $uv \in \delta(W)$  (say  $u \in W_t$  and  $v \in W_h$ ) and  $uv \in \delta(Y)$  (say  $v \in Y_t$  and  $u \in Y_h$ ). Hence, there is a type I setpair  $W \in \mathcal{L}$  such that  $\delta(W)$  has size at most 12, and so there is an edge  $e \in \delta(W)$  with  $x_e \geq \frac{1}{12}$ .

(Type II) Suppose that there are at least  $\frac{m}{6}$  type II setpairs. Let  $Y^1, \dots, Y^q$  be the minimal nodes in  $\mathcal{P}$  (those with no children); clearly, these nodes are pairwise incomparable. Each  $Y^i$  and its ancestors forms a (directed) tree (since  $\mathcal{P}$  is diamond-free). Each node of  $\mathcal{P}$  is in at most  $\frac{2}{\epsilon}$  of these trees, by Lemma 4.4. Focus on one of the trees and note that the number of leaf nodes (maximal nodes of  $\mathcal{P}$ ) of the tree is greater than or equal to the number of type II nodes in the tree. Summing over all the trees, we see that the total number of leaf nodes (maximal nodes of  $\mathcal{P}$ ) is at least  $\frac{\epsilon}{2}$  times the number of type II nodes in  $\mathcal{P}$ , hence,  $\mathcal{P}$  has at least  $\frac{m\epsilon}{12}$  maximal nodes; clearly, these nodes are pairwise incomparable. Then the edge sets  $\delta(W)$ , where  $W$  is a maximal node of  $\mathcal{P}$ , are quasidisjoint, so there is an edge  $e$  with  $x_e \geq \frac{\epsilon}{24}$ .

(Type III) Finally, suppose that there are at least  $\frac{m}{6}$  type III setpairs. Consider all of the type III setpairs  $W^1, W^2, \dots, W^q$  and their unique parents  $Y^1, Y^2, \dots, Y^q$ . Observe that  $f(W^i) + 1 \leq f(Y^i)$ , since  $|W_t^i \cup W_h^i| < |Y_t^i \cup Y_h^i|$ . Moreover, if  $W^i$  and  $W^j$  (where  $1 \leq i < j \leq q$ ) have distinct parents  $Y^i$  and  $Y^j$ , then the edge sets  $\delta(Y^i) \setminus \delta(W^i)$  and  $\delta(Y^j) \setminus \delta(W^j)$  are quasidisjoint. To see this, consider an edge  $uv \in \delta(Y^i) \setminus \delta(W^i)$  with  $u \in Y_t^i = W_t^i$  and  $v \in Y_h^i - W_h^i$ . Now, every proper ancestor  $Z$  of  $W^i$  has  $v \in Z_h$ , so if  $W^j$  is an ancestor of  $W^i$ , then we cannot have  $uv \in \delta(Y^j) \setminus \delta(W^j)$ ,  $u \in Y_t^j = W_t^j$ , and  $v \in Y_h^j - W_h^j$ .

Consequently, by Lemma 4.5(b), any edge occurs in at most  $\frac{4}{\epsilon}$  of the edge sets  $\delta(Y^i) \setminus \delta(W^i)$ , and moreover,  $x(\delta(Y^i) \setminus \delta(W^i)) \geq 1$ , for each  $i = 1, \dots, q$ . Hence, there is an edge  $e$  with  $x_e \geq \frac{\epsilon}{24}$ .

This concludes the proof of Theorem 4.2 for undirected graphs: any nonzero basic solution of (LP-VC) has an edge of value  $\Omega(\sqrt{\epsilon/n})$ .

The proof extends to directed graphs via Theorem 4.3. We partition the noncrossing family of setpairs  $\mathcal{L}$  into two subfamilies  $\mathcal{L}_1$  and  $\mathcal{L}_2$ , depending

on whether or not a setpair  $W \in \mathcal{L}$  has  $|W_h| \geq |W_t|$ . Suppose that  $|\mathcal{L}_1| \geq |\mathcal{L}_2|$ ; otherwise, we use a symmetric argument. Then we apply the arguments for undirected graphs to  $\mathcal{L}_1$  (partitioning it into types I, II, III, IV, etc.). We obtain similar lower bounds on  $\max\{x_e\}$ , the main difference being that we lose a factor of two in the lower bound (since we have  $|\mathcal{L}_1| \geq \frac{m}{2}$  rather than  $|\mathcal{L}| = m$ ). Hence, Theorem 4.2 holds for directed graphs. ■

Theorem 4.1 follows from Theorem 4.2: for  $k \leq (1-\epsilon)n$ , iterative rounding gives an  $O(\sqrt{n/\epsilon})$  approximation algorithm for the (directed or undirected)  $k$ -vertex connectivity problem.

Is it possible to improve substantially on our analysis of the iterative rounding method for the  $k$ -vertex connectivity problem? This is not clear at present, since our approximation guarantee holds for directed graphs for large values of  $k$  (say  $n/2 < k < n - \Omega(1)$ ), and for this case, the previous best approximation guarantee was  $\Omega(k)$ . Moreover, there is an example of the undirected problem such that there is a basic solution  $\mathbf{x}$  of (LP-VC) such that each edge  $e$  has  $x_e \leq \frac{1}{\Omega(\sqrt{k})}$ . Hence, in Theorem 4.2, the lower bound on the maximum value of an edge cannot be improved beyond  $\frac{1}{\Omega(\sqrt{k})}$ . We have no new lower bounds for the integrality ratio of (LP-VC).

### 5. An Approximation Algorithm for the Element Connectivity Problem

This section proves a 2-approximation guarantee for a generalization of the element connectivity problem. A symmetric, non-negative, integral function  $\psi$  on the subsets of a groundset  $U$  is called *weakly supermodular* if  $\psi(U) = 0$ , and for every two subsets  $X$  and  $Z$ , we have either  $\psi(X) + \psi(Z) \leq \psi(X \cap Z) + \psi(X \cup Z)$  or  $\psi(X) + \psi(Z) \leq \psi(\overline{X} \cap Z) + \psi(\overline{X} \cup Z)$ , where  $\overline{X} = U \setminus X$  ( $\psi$  is symmetric if  $\psi(X) = \psi(\overline{X}), \forall X \subseteq U$ ). Recall that  $T \subseteq V$  is given as the set of terminal vertices of an undirected graph  $G = (V, E)$ , and each edge  $e$  has a nonnegative cost  $c_e$ . In the *generalized element connectivity problem*, rather than specifying the connectivity requirements by a function  $r: (T \times T) \rightarrow \mathbb{Z}_+$ , we use a weakly supermodular function on the set of terminals  $g': 2^T \rightarrow \mathbb{Z}_+$  (we assume that  $g'$  is symmetric, it assigns a nonnegative integer value to each subset of  $T$ , and  $g'(T) = g'(\emptyset) = 0$ ). For the integer program (IP), the requirement function  $f$  on the setpairs  $W \in \mathcal{S}$  is defined via  $g'$  as follows:

$$f(W) = \begin{cases} \max\{0, g'(W_h \cap T) - |V \setminus (W_t \cup W_h)|\}, & \text{if } T \subseteq W_t \cup W_h; \\ 0, & \text{otherwise.} \end{cases}$$

Note that if a setpair  $W$  has a positive requirement, then all the terminals are in  $W_t \cup W_h$ .

Let (LP-EC) denote the linear programming relaxation of this integer program. Theorem 5.5 shows a key property of (LP-EC), namely, there exists an edge of value at least  $\frac{1}{2}$  in any non-zero basic solution. If a polynomial-time strong separation oracle for (LP-EC) is available, then a basic optimal solution to (LP-EC) (if it exists) can be found in polynomial time. Thus the next result follows from the iterative rounding method.

**Theorem 5.1.** *Consider the generalized element connectivity problem, and suppose that a strong separation oracle is available (for the linear programming relaxation). Then there is a polynomial-time algorithm that given an instance of the problem, finds a solution of cost at most twice the optimal cost.* ■

For the special case of the element connectivity problem, a strong separation oracle is available, and thus we obtain an efficient algorithm with an approximation guarantee of 2; this matches the earlier result of Fleischer, Jain, and Williamson [7].

**Lemma 5.2.** *If the function  $g'$  is weakly supermodular, then the requirement function  $f$  for the generalized element connectivity problem is skew bisupermodular.*

**Proof.** Define a function  $g(\cdot)$  on the setpairs  $W \in \mathcal{S}$  by  $g(W) = g'(W_h \cap T)$ . We claim that the function  $g$  is skew bisupermodular. This follows from the term by term correspondence between the skew bisupermodular inequalities for  $g$  and the weakly supermodular inequalities for  $g'$ . (To see this in detail, consider overlapping setpairs  $W$  and  $Y$  with  $g(W) > 0, g(Y) > 0$ . Clearly, for both  $W$  and  $Y$ ,  $T$  is contained in the union of the head and the tail. Then both  $W \otimes Y$  and  $W \oplus Y$  also have this property. Let  $X = W_h \cap T$  and let  $Z = Y_h \cap T$ . Suppose that one of the weakly supermodular inequalities holds for  $X$  and  $Z$ , say,  $g'(X) + g'(Z) \leq g'(X \cap Z) + g'(X \cup Z)$ . Note that  $g'(X) = g(W)$ ,  $g'(Z) = g(Y)$ ,  $g'(X \cap Z) = g(W \otimes Y)$ , and  $g'(X \cup Z) = g(W \oplus Y)$ . Hence, the skew bisupermodular inequality  $g(W) + g(Y) \leq g(W \otimes Y) + g(W \oplus Y)$  holds for  $W$  and  $Y$ .)

Now, consider the function  $f(\cdot)$ . We may write  $f(\cdot)$  as

$$f(W) = \begin{cases} \max\{0, g(W) - p(W)\}, & \text{if } T \subseteq W_t \cup W_h; \\ 0, & \text{otherwise;} \end{cases}$$

where  $p(W)$  denotes  $|V \setminus (W_t \cup W_h)|$  for each setpair  $W$ . Recall from Section 4 that for any two setpairs  $W, Y$  we have  $p(W) + p(Y) = p(W \otimes Y) + p(W \oplus Y)$ ,

and  $p(W) + p(Y) = p(\overline{W} \otimes Y) + p(\overline{W} \oplus Y)$ . Thus it can be seen that  $f(\cdot)$  is skew bisupermodular.  $\blacksquare$

Now, consider the special case of the element connectivity problem. For all  $i, j \in T$ , there is a requirement for  $r_{ij}$  element disjoint paths between  $i$  and  $j$ . Define the function  $g'(\cdot)$  on  $T$  as follows. Let  $g'(S) = \max_{i \in S, j \notin S} \{r_{ij}\}$  for any subset  $S$  of  $T$ . (Note that  $g'$  is symmetric, nonnegative, integral, and  $g'(T) = g'(\emptyset) = 0$ .) Frank [3, Proposition 5.4] (also Goemans et al [8]) showed that  $g'$  is weakly supermodular. Define the requirement function  $f(\cdot)$  on the setpairs  $W \in \mathcal{S}$  as in the generalized problem. It is easily seen that the integer program (IP) models the element connectivity problem (for more details see [12]). Lemma 5.2 applies and shows that  $f$  is a skew bisupermodular function. Moreover, for this special case, a strong separation oracle for (LP-EC) is available using standard network flow techniques, hence, a basic optimal solution to (LP-EC) (if it exists) can be found in polynomial time. Thus Theorem 5.1 gives an (efficient) 2-approximation algorithm for the element connectivity problem (without further assumptions).

To prove Theorem 5.5 (there exists an edge of value at least  $\frac{1}{2}$  in any non-zero basic solution of (LP-EC)), we need to develop some preliminaries and a key lemma. Based on this, the rest of the proof becomes an easy extension of the arguments in Jain's proof of [11, Lemma 4.6].

Let  $\mathbf{x}$  be a non-zero basic solution of (LP-EC). We may assume that each edge  $e \in E$  has  $x_e > 0$ , since any other edges may be discarded. In addition, we may assume that each edge  $e \in E$  has  $x_e < 1$ , otherwise we are done. Let  $\mathcal{L}$  be the corresponding non-overlapping family of setpairs of Theorem 3.3. Since  $E$  is the support of a basic solution,  $|E| = |\mathcal{L}|$ . Let  $m$  denote  $|E|$ . Let  $\tau$  be an arbitrarily chosen terminal vertex in  $T$ . For each setpair in  $\mathcal{L}$ , we assume that the head contains  $\tau$ . Otherwise, if  $\tau$  is in the tail of  $W \in \mathcal{L}$ , then we replace  $W$  by  $\overline{W}$  and this preserves all the properties of  $\mathcal{L}$  in Theorem 3.3. Then the tails of the setpairs in  $\mathcal{L}$  form a laminar family. This holds because the setpairs in  $\mathcal{L}$  are pairwise non-overlapping, and the heads of any two setpairs in  $\mathcal{L}$  intersect (since  $\tau$  is in every head). Moreover, each setpair  $W \in \mathcal{L}$  has a terminal in its tail (since  $f(W) > 0$  implies  $g'(W_h \cap T) > 0$ , so  $W_h \cap T \subset T$ ). We may also view  $\mathcal{L}$  as a forest of rooted trees. Call  $W \in \mathcal{L}$  the *parent* of  $Y \in \mathcal{L}$  (and  $Y$  a *child* of  $W$ ) if  $W_t \supseteq Y_t$  and any other setpair  $Z \in \mathcal{L}$  with  $Z_t \supseteq Y_t$  either has  $Z_t \supset W_t$  or has  $Z_t = W_t$  and  $Z_h \subset W_h$ . We may refer to the setpairs in  $\mathcal{L}$  as *nodes* of the forest. We say that a setpair  $Y \in \mathcal{L}$  is *smaller* than a setpair  $W \in \mathcal{L}$  if  $Y \succeq W$ , i.e.,  $Y_t \subseteq W_t$  and  $Y$  is a descendant of  $W$  in the forest.

**Lemma 5.3.** *Let  $Y = (Y_t, Y_h)$  and  $Z = (Z_t, Z_h)$  be setpairs in  $\mathcal{L}$ . If  $Y$  and  $Z$  are tail disjoint, then  $Y_t \subseteq Z_h$  and  $Z_t \subseteq Y_h$ .*

**Proof.** Note that  $Y_t$  and  $Z_h$  have a terminal in common, since  $Y_t$  has at least one terminal, and  $Z_t \cup Z_h$  contains all the terminals. Similarly,  $Z_t$  and  $Y_h$  have a terminal in common. Suppose that  $Y_t$  is not a subset of  $Z_h$ . Then it can be seen that  $Y_t$  and  $Z_h$  intersect properly (since  $\tau \in Z_h \setminus Y_t$  and there is a vertex in  $Y_t \setminus Z_h$ ). Hence,  $\overline{Z}$  crosses  $Y$ , and so  $Y$  and  $Z$  overlap. This contradiction shows that  $Y_t \subseteq Z_h$ . Similarly, we have  $Z_t \subseteq Y_h$ .  $\blacksquare$

In the rest of this section, we take the edges in  $E$  to be bidirected. That is, we replace every undirected edge  $\{p, q\}$  in  $E$  by a pair of directed arcs  $pq$  and  $qp$ . For a setpair  $W = (W_t, W_h)$ , let  $\delta(W)$  denote the set of arcs  $pq$  with  $p \in W_t$  and  $q \in W_h$ . (Though we use the same notation  $\delta()$  for undirected edges and directed arcs, the context will resolve any ambiguity.) The other notation,  $\mathbf{x}$ ,  $\chi_W$ , etc. remains the same. An arc  $pq$  is called *good* if there is a setpair  $W \in \mathcal{L}$  such that  $pq$  is in  $\delta(W)$ . Note that, in the undirected setting, every edge  $e$  is in  $\delta(W)$  for some  $W \in \mathcal{L}$ , but in the directed setting, there may exist arcs that are not good.

The proof of the main result hinges on assigning *token arcs* to the setpairs in  $\mathcal{L}$ . Here is a brief sketch; the details are given in the proof of Theorem 5.5. We have  $|\mathcal{L}| = |E| = m$  and thus a total of  $2m$  arcs. If a basic solution  $\mathbf{x}$  of (LP-EC) has  $x_e < \frac{1}{2}$  for each edge  $e$ , then we will be able to distribute the token arcs among the setpairs in  $\mathcal{L}$  such that each gets at least two token arcs and some setpair gets at least three token arcs. Thus, we end up with at least  $2|\mathcal{L}| + 1 = 2m + 1$  arcs. This contradiction shows that our assumption on  $\max_e \{x_e\}$  is false. We use the following two rules to distribute the token arcs such that no arc is assigned to two different setpairs.

**Rule 1.** If  $\alpha\beta$  is a good arc, then we assign it to the smallest setpair  $W \in \mathcal{L}$  such that  $\alpha\beta \in \delta(W)$ .

**Rule 2.** If  $\alpha\beta$  is not a good arc, then we assign it to the smallest setpair  $W \in \mathcal{L}$  such that  $\beta \in W_t$  and  $\alpha \notin W_h$ .

(Rule 2 is essential in the sense that the analysis fails if we insist on assigning arcs  $pq$  only to those setpairs  $W$  such that  $\{p, q\} \in \delta(W)$ .)

Given two setpairs  $W$  and  $Y$ , let  $\delta(W) \nabla \delta(Y)$  denote the symmetric difference of  $\delta(W)$  and  $\delta(Y)$ . Given  $S^1, S^2, \dots, S^\ell$ , a collection of mutually disjoint vertex sets, we denote by  $\gamma(S^1, S^2, \dots, S^\ell)$  the set of arcs  $pq$  such that  $p$  and  $q$  are in different sets  $S^i$  and  $S^j$ , where  $1 \leq i, j \leq \ell$ .




**Lemma 5.4.** *Let  $W$  be a setpair in the forest  $\mathcal{L}$ , and let its children be the setpairs  $Y^1, Y^2, \dots, Y^\ell$ . Then for every arc  $pq$  in*

$$(\delta(W) \nabla (\delta(Y^1) \cup \dots \cup \delta(Y^\ell))) \setminus \gamma(Y_t^1, \dots, Y_t^\ell)$$

*either  $pq$  or  $qp$  is assigned to  $W$  by Rule 1 or Rule 2.*

**Proof.** Note that the lemma applies to all arcs in the symmetric difference of  $\delta(W)$  and  $\bigcup_{i=1}^\ell \delta(Y^i)$  except for arcs whose two end-vertices are in the tails of different children. Figure 1 illustrates the arguments that follow, for the particular case of  $\ell=2$ . [Note, for example, that the arc  $a_4$  is in  $\gamma(Y_t^1, Y_t^2)$ .]



**Figure 1.** An illustration of the proof of Lemma 5.4.

We have two cases to deal with.

(a)  $pq \in \delta(W) \setminus \bigcup_{i=1}^\ell \delta(Y^i)$ .

Now  $q \in W_h \subseteq Y_h^i$ , for  $i = 1, \dots, \ell$ . However, by assumption,  $p \in W_t \setminus \bigcup_{i=1}^\ell Y_t^i$  and so  $W$  is the smallest setpair in  $\mathcal{L}$  such that  $p$  is in the tail. Thus,  $W$  is the smallest setpair in  $\mathcal{L}$  with  $pq \in \delta(W)$ , and hence  $pq$  is assigned to  $W$  by Rule 1. [The arc  $a_1$  in Figure 1 is an example.]

(b)  $pq \in \delta(Y^i) \setminus \delta(W)$ , for some  $i \in \{1, \dots, \ell\}$ , and  $pq \notin \gamma(Y_t^1, \dots, Y_t^\ell)$ .

Note that  $p \in Y_t^i \subseteq W_t$  and  $q \notin W_h$ . We have two subcases.

(i)  $q \in W_t$ .

First, we show that  $qp$  is not a good arc. Clearly,  $q \notin Y_t^i$  as  $pq \in \delta(Y^i)$ . For any other child  $Y^j$ , observe that  $q \notin Y_t^j$  as  $pq \notin \gamma(Y_t^1, \dots, Y_t^\ell)$ . Hence,  $q \in W_t \setminus \cup_{j=1}^\ell Y_t^j$ . Then  $W$  is the smallest setpair in  $\mathcal{L}$  whose tail contains  $q$ , and there is no setpair  $Z \in \mathcal{L}$  such that  $qp \in \delta(Z)$  (otherwise,  $p \in Z_h \cap W_t$  and  $q \in W_t \subseteq Z_t$ , which is impossible). Thus  $qp$  is not a good arc. Moreover,  $p \notin W_h$  (since  $p \in W_t$ ), and so Rule 2 assigns  $qp$  to  $W$ . [The arc  $a_2$  in Figure 1 is an example.]

(ii)  $q \notin W_t$ .

We first show that  $qp$  is not a good arc. Suppose  $Z \in \mathcal{L}$  is a setpair such that  $qp \in \delta(Z)$ . Note that  $W_t$  and  $Z_t$  are disjoint, because  $p \in W_t \setminus Z_t$  and  $q \in Z_t \setminus W_t$ . Then by Lemma 5.3  $Z_t \subseteq W_h$ . This gives a contradiction (since  $q \in Z_t$ ,  $q \notin W_h$ ). Therefore,  $qp$  is not a good arc and Rule 2 applies to  $qp$ . Rule 2 assigns  $qp$  to  $W$  since  $p \in W_t$ ,  $q \notin W_h$ , and for any smaller setpair  $X \in \mathcal{L}$  with  $p \in X_t$  we have  $q \in X_h$  (since  $p \in Y_t^i$  and  $q \in Y_h^i$ ). [The arc  $a_3$  in Figure 1 is an example.]  $\blacksquare$

The next result is proved by contradiction, and uses similar arguments to Jain's proof of [11, Lemma 4.6]. We include the proof, for the sake of completeness.

**Theorem 5.5.** *Any non-zero basic solution of (LP-EC) has an edge of value at least  $\frac{1}{2}$ .*

**Proof.** Let  $\mathbf{x} \neq \mathbf{0}$  be a basic solution of (LP-EC) such that  $x_e < \frac{1}{2}$  for each edge  $e$ . Then the following claim implies that we end up with at least  $2|\mathcal{L}|+1$  token arcs. This is a contradiction. Hence,  $\max_e \{x_e\} \geq \frac{1}{2}$ .

Define the *corequirement* of an edge  $e$ , denoted  $\alpha(e)$ , to be  $\frac{1}{2} - x_e > 0$ , and the *corequirement* of a setpair  $W$ , denoted  $\alpha(W)$ , to be  $\sum_{e \in \delta(W)} \alpha(e)$ . For a tight setpair  $W$ , note that  $\alpha(W) = \frac{1}{2}|\delta(W)| - f(W) > 0$ , hence, if  $\alpha(W) = \frac{1}{2}$ , then  $|\delta(W)|$  is odd. Also, note that if  $\delta(W) \subseteq \delta(Y^1) \cup \dots \cup \delta(Y^\ell)$ , then  $\alpha(W) = \sum_{e \in \delta(W)} \alpha(e) \leq \alpha(Y^1) + \dots + \alpha(Y^\ell)$ . (Observe that corequirements are not defined for arcs.)

**Claim.** *Suppose that  $x_e < \frac{1}{2}, \forall e \in E$ . Then the arcs may be redistributed to the nodes in the forest  $\mathcal{L}$  such that for each rooted subtree of  $\mathcal{L}$  each node gets at least two token arcs and the root gets at least three token arcs. Moreover, a root with corequirement  $\neq \frac{1}{2}$  gets at least four token arcs.*

We prove the claim by induction. For the base case take a leaf node  $W \in \mathcal{L}$ . Then  $\delta(W)$  has at least 3 arcs. By Rule 1, all these arcs are assigned to  $W$ . If  $W$  gets exactly 3 token arcs, then note that  $f(W) = 1$  and  $|\delta(W)| = 3$ , so we have  $\alpha(W) = \frac{1}{2}$ . Thus  $W$  satisfies the induction hypothesis.

For the induction step, consider a subtree rooted at a node  $W \in \mathcal{L}$ . We have four cases.

(a)  $W$  has at least four children.

By the induction hypothesis, each child has at least 3 token arcs. We reassign one token arc from each child to  $W$ . Hence,  $W$  gets at least 4 token arcs.

(b)  $W$  has three children.

Call the children  $X$ ,  $Y$  and  $Z$ . If one of the children has at least 4 token arcs, then we can reassign the token arcs from  $X$ ,  $Y$  and  $Z$  to  $W$  so that  $W$  has at least 4 token arcs and each of  $X$ ,  $Y$  and  $Z$  is left with at least 2 token arcs.

Otherwise, by the induction hypothesis, each of  $X$ ,  $Y$  and  $Z$  has 3 token arcs, so each has corequirement  $\frac{1}{2}$  and each of  $|\delta(X)|$ ,  $|\delta(Y)|$  and  $|\delta(Z)|$  is odd. We reassign one token arc from each of  $X$ ,  $Y$  and  $Z$  to  $W$ . Now focus on the symmetric difference of  $\delta(W)$  and  $\delta(X) \cup \delta(Y) \cup \delta(Z)$ . By the linear independence of  $\chi_W$ ,  $\chi_X$ ,  $\chi_Y$  and  $\chi_Z$ , there is at least one arc  $pq$  in the symmetric difference. If possible, choose  $pq \notin \gamma(X_t, Y_t, Z_t)$ . Then by Lemma 5.4 either  $pq$  or  $qp$  is assigned to  $W$  as a token arc. Hence,  $W$  ends up with at least 4 token arcs. Otherwise, every arc in the symmetric difference is in  $\gamma(X_t, Y_t, Z_t)$ . Therefore  $\delta(W) \subset \delta(X) \cup \delta(Y) \cup \delta(Z)$ , and for each arc  $pq$  in  $(\delta(X) \cup \delta(Y) \cup \delta(Z)) \setminus \delta(W)$  the arc  $qp$  is also in the same set (by Lemma 5.3). Hence,  $|\delta(W)|$  is odd, and so the corequirement  $\alpha(W)$  is a *semi integer* (i.e., an odd integer multiple of  $\frac{1}{2}$ ). Moreover,  $\alpha(W) < \alpha(X) + \alpha(Y) + \alpha(Z) = \frac{3}{2}$ . Hence,  $\alpha(W) = \frac{1}{2}$ , and so  $W$  satisfies the induction hypothesis.

(c)  $W$  has two children.

Call the children  $Y$  and  $Z$ . If we can assign 4 token arcs to  $W$  by reassigning token arcs from  $Y$  and  $Z$ , and by applying Lemma 5.4 to  $(\delta(W) \nabla (\delta(Y) \cup \delta(Z))) \setminus \gamma(Y_t, Z_t)$ , then we are done. Otherwise, we are left with two subcases.

(i) Each of  $Y$  and  $Z$  has 3 token arcs and a corequirement of  $\frac{1}{2}$ , and there is one arc  $pq$  in  $(\delta(W) \nabla (\delta(Y) \cup \delta(Z))) \setminus \gamma(Y_t, Z_t)$ . Then  $W$  gets 3 token arcs, one each by reassigning arcs from  $Y$  and  $Z$  and one by Lemma 5.4. Both  $|\delta(Y)|$  and  $|\delta(Z)|$  are odd (since  $\alpha(Y) = \alpha(Z) = \frac{1}{2}$ ) and the arcs (if any) in  $\gamma(Y_t, Z_t)$  occur in pairs  $uv$  and  $vu$  (by Lemma 5.3). Therefore,  $|\delta(W)|$  is odd and  $\alpha(W)$  is a semi integer. Moreover,  $0 < \alpha(W) \leq \alpha(Y) + \alpha(Z) + \alpha(\{p, q\}) < \frac{3}{2}$ . Hence,  $\alpha(W) = \frac{1}{2}$ , and so  $W$  satisfies the induction hypothesis.

(ii) At least one of the children, say  $Z$ , has 3 token arcs and a corequirement of  $\frac{1}{2}$ , and there are no arcs in  $(\delta(W) \nabla (\delta(Y) \cup \delta(Z))) \setminus \gamma(Y_t, Z_t)$ . We will prove by contradiction that this case cannot occur. First, note that  $|\delta(W)|$  and  $|\delta(Y)|$  have opposite parity (one is odd and the other is even),

because  $|\delta(Z)|$  is odd and the arcs in the symmetric difference of  $\delta(W)$  and  $\delta(Y) \cup \delta(Z)$  occur in pairs  $uv$  and  $vu$ . Furthermore, note that  $\chi_W, \chi_Y$  and  $\chi_Z$  are linearly independent. It follows that there is an arc in  $\gamma(Y_t, Z_t)$  and there is an arc in  $\delta(Z) \cap \delta(W)$ . Hence,  $\alpha(W) < \alpha(Y) + \alpha(Z)$  (due to the arc in  $\gamma(Y_t, Z_t)$ ) and  $\alpha(Y) < \alpha(W) + \alpha(Z)$  (due to the arc in  $\delta(Z) \cap \delta(W)$ ). Therefore,  $\alpha(Y) - \frac{1}{2} < \alpha(W) < \alpha(Y) + \frac{1}{2}$ , implying that  $\alpha(Y) = \alpha(W)$  (since  $\alpha(W)$  is an integer multiple of  $\frac{1}{2}$ ). Then  $|\delta(W)|$  and  $|\delta(Y)|$  have the same parity. This gives the desired contradiction.

(d)  $W$  has one child.

Call the child  $Y$ . If the symmetric difference of  $\delta(W)$  and  $\delta(Y)$  has exactly 2 arcs and  $\alpha(Y) = \frac{1}{2}$ , then  $W$  gets 3 token arcs (two by Lemma 5.4 and one reassigned from  $Y$ ). In this case, note that  $f(W) = f(Y)$  and  $|\delta(W)| = |\delta(Y)|$ . As a result,  $\alpha(W) = \alpha(Y) = \frac{1}{2}$ . Otherwise,  $W$  gets at least 4 token arcs. In either case,  $W$  satisfies the induction hypothesis. This completes the proof of the claim.

The theorem follows. ■

Theorem 5.1 follows from Theorem 5.5 by applying the iterative rounding method to (LP-EC). Note that if  $f$  is a (non-negative, integral, symmetric) skew bisupermodular function on  $\mathcal{S}$  and  $e$  is any edge, then the function  $f'$  is also skew bisupermodular, where  $f'(W) = \max\{0, f(W) - 1\}$ , if  $e \in \delta(W)$ , and  $f'(W) = f(W)$ , otherwise.

## References

- [1] J. CHERIYAN and S. VEMPALA: Edge coverings of setpairs and the iterative rounding method, submitted for journal publication, September 2001. Preliminary version in the *Proceedings of the 8th International Integer Programming and Combinatorial Optimization Conference*, Utrecht, The Netherlands, pp. 30–44, 2001.
- [2] J. CHERIYAN, S. VEMPALA and A. VETTA: An approximation algorithm for the minimum-cost  $k$ -vertex connected subgraph, *SIAM Journal on Computing* **32** (2003), 1050–1055.
- [3] A. FRANK: Augmenting graphs to meet edge-connectivity requirements, *SIAM J. Discrete Math.* **5** (1992), 25–53. Preliminary version in *Proc. 31st IEEE FOCS*, pp. 708–718, 1990.
- [4] A. FRANK: Connectivity augmentation problems in network design, in *Mathematical Programming: State of the Art 1994*, (Eds. J. R. Birge and K. G. Murty), pp. 34–63, The University of Michigan, Ann Arbor, MI, 1994.
- [5] A. FRANK and T. JORDAN: Minimal edge-coverings of pairs of sets, *Journal of Combinatorial Theory, Series B* **65** (1995), 73–110.
- [6] A. FRANK and É. TARDOS: An application of submodular flows, *Linear Algebra and its Applications* **114/115** (1989), 329–348.

- [7] L. FLEISCHER, K. JAIN and D. P. WILLIAMSON: An iterative rounding 2-approximation algorithm for the element connectivity problem, to appear in the *Journal of Computer and System Sciences*. Preliminary version in *Proc. 42nd IEEE FOCS*, pp. 339–347, 2001.
- [8] M. GOEMANS, A. GOLDBERG, S. PLOTKIN, D. SHMOYS, É. TARDOS and D. WILLIAMSON: Improved approximation algorithms for network design problems, *Proc. 5th ACM-SIAM SODA*, pp. 223–232, 1994.
- [9] M. GOEMANS and D. WILLIAMSON: A general approximation technique for constrained forest problems, *SIAM Journal on Computing* **24** (1995), 296–317.
- [10] M. GRÖTSCHEL, L. LOVÁSZ and A. SCHRIJVER: *Geometric Algorithms and Combinatorial Optimization*, Springer-Verlag, New York, 1988.
- [11] K. JAIN: A factor 2 approximation algorithm for the generalized Steiner network problem, *Combinatorica* **21** (2001), 39–60. Preliminary version in *Proc. 39th IEEE FOCS*, 1998.
- [12] K. JAIN, I. MANDOIU, V. VAZIRANI and D. WILLIAMSON: A primal-dual schema based approximation algorithm for the element connectivity problem, *Journal of Algorithms* **45** (2001), 1–15. Preliminary version in *Proc. 10th ACM-SIAM SODA*, pp. 484–489, 1999.
- [13] G. KORTSARZ and Z. NUTOV: Approximating  $k$ -node connected subgraphs via critical graphs, *Proc. 36th ACM STOC*, June 2004.
- [14] G. KORTSARZ and Z. NUTOV: Approximating node connectivity problems via set covers, *Algorithmica* **37** (2003), 75–92. Preliminary version in *APPROX, Approximation algorithms for combinatorial optimization*, Springer, LNCS **1913**, pp. 194–205, 2000.
- [15] R. RAVI and D. P. WILLIAMSON: An approximation algorithm for minimum-cost vertex-connectivity problems, *Algorithmica* **18** (1997), 21–43. Preliminary version in *Proc. 6th ACM-SIAM SODA*, pp. 332–341, 1995.  
Erratum, *Algorithmica* **34** (2002), 98–107.
- [16] É. TARDOS: A strongly polynomial algorithm to solve combinatorial linear programs, *Operations Research* **34** (1986), 250–256.

Joseph Cheriyan

*Department of Combinatorics and Optimization*  
*University of Waterloo*  
 200 University Ave. West  
 Waterloo, ON, N2L 3G1  
 Canada  
 jcheriyan@math.uwaterloo.ca

Santosh Vempala

*Department of Mathematics*  
*M.I.T.*  
 Cambridge, MA 02139  
 USA  
 vempala@math.mit.edu

Adrian Vetta

*Department of Mathematics and Statistics*  
*McGill University*  
 Montreal, Quebec, H3A 2A7  
 Canada  
 vetta@math.mcgill.ca