

An equation

$$\gamma_1(x) = P(x) - \gamma_1(x)(1 - sx\partial_x)\gamma_1(x)$$

Important special cases

$$\begin{aligned} \gamma_1(x) &= x - \gamma_1(x)(1 - 2x\partial_x)\gamma_1(x) \\ 2\gamma_1(x) &= \left(\frac{x}{3} + \frac{x^2}{4} \right) - \gamma_1(x)(1 - x\partial_x)\gamma_1(x) \\ 2\gamma_1(x) &= \left(\frac{x}{3} + \frac{x^2}{4} + (-0.0312 + 0.06037)x^3 + \right. \\ &\quad \left. (-0.6755 + 0.05074)x^4 \right) \\ &\quad - \gamma_1(x)(1 - x\partial_x)\gamma_1(x) \end{aligned}$$

$$\begin{aligned} \gamma_1^+(x) &= P^+(x) - \gamma_1^+(x)^2 - (\gamma_1^+(x) - 2\gamma_1^-(x))x\partial_x\gamma_1^+(x) \\ \gamma_1^-(x) &= P^-(x) - \gamma_1^-(x)^2 - (\gamma_1^+(x) - 2\gamma_1^-(x))x\partial_x\gamma_1^-(x) \end{aligned}$$

0-1

Getting there

In the context of renormalization Hopf algebras consider

$$X(x) = \mathbb{I} - \sum_{k \geq 1} x^k p(k) B_+^k (X(x) Q(x)^k)$$

where $Q(x) = X(x)^{-s}$ with $s > 0$ an integer. Associate with each B_+ a

$$F^k(\rho).$$

Write the combination $(X \mapsto G, B_+^k \mapsto F^k, \rho \text{ marks the insertion place})$ as $G(x, L) = \sum \gamma_k(x) L^k$ with $\gamma_k(x) = \sum_{j \geq k} \gamma_{k,j} x^j$.

Systems of equation are similar but messier.

0-2

The $\gamma_k(x)$ recursion

We know from Connes and Kreimer [2] that if

$$\sigma_1 = \partial_L \phi_R(S \star Y)|_{L=0} \quad \text{and} \quad \sigma_n = \frac{1}{n!} m^{n-1} \underbrace{(\sigma_1 \otimes \cdots \otimes \sigma_1)}_n \Delta^{n-1}$$

then

$$\gamma_k(x) = \sigma_k(X(x)).$$

But σ_1 only sees the linear part of the Hopf algebra so we can use $\Delta_{\text{lin}} = (P_{\text{lin}} \otimes \cdots \otimes P_{\text{lin}}) \Delta^{n-1}$ in place of Δ where P_{lin} projects onto the linear part of the Hopf algebra.

Calculate

$$\Delta_{\text{lin}} X = P_{\text{lin}} X \otimes P_{\text{lin}} X + P_{\text{lin}} Q \otimes x\partial_x X.$$

So

$$\gamma_k(x) = \frac{1}{k} \gamma_1(x)(1 - sx\partial_x)\gamma_{k-1}(x),$$

0-3

The γ_1 recursion

Rewrite the (analytic) Dyson-Schwinger equation

$$\gamma \cdot L = \sum p(k)x^k(1 + \gamma \cdot \partial_{-\rho})^{sk+1}(1 - e^{-L\rho})F^k(\rho) \Big|_{\rho=0}$$

where $\gamma \cdot U = \sum \gamma_k U^k$.

Take an L derivative and set $L = 0$ to get

$$\gamma_1 = \sum p(k)x^k(1 + \gamma \cdot \partial_{-\rho})^{sk+1}\rho F^k(\rho) \Big|_{\rho=0}$$

Assume $\rho F^k(\rho) = r_k/(1 - \rho)$ and take two L derivatives to get

$$2\gamma_2 = - \sum_k p(k)x^k(1 + \gamma \cdot \partial_{-\rho})^{sk+1}r_k \frac{\rho}{1 - \rho} \Big|_{\rho=0} = -\gamma_1 + \sum x^k p(k)r_k.$$

Write $P(x) = \sum x^k p(k)r_k$ and use the other recursion:

$$\gamma_1 = P(x) - \gamma_1(1 - sx\partial_x)\gamma_1.$$

0-4

Solved

Broadhurst and Kreimer [1] solved this Dyson-Schwinger equation by clever rearranging and recognizing the resulting asymptotic expansion. Today Maple can solve it.

$$\gamma_1(x) = x - \gamma_1(x)(1 - 2x\partial_x)\gamma_1(x)$$

gives

$$\exp\left(\frac{(1 + \gamma_1(x))^2}{2x}\right) \sqrt{-x} + \operatorname{erf}\left(\frac{1 + \gamma_1(x)}{\sqrt{-2x}}\right) \frac{\sqrt{\pi}}{\sqrt{2}} = C$$

0-6

Where it all began

Broadhurst and Kreimer [1]; a bit of massless Yukawa theory.

$$X(x) = \mathbb{I} - xB_+ \left(\frac{1}{X(x)} \right),$$

$$F(\rho) = \frac{1}{q^2} \int d^4 k \frac{k \cdot q}{(k^2)^{1+\rho}(k+q)^2} - \dots \Big|_{q^2=\mu^2}.$$

Combine to get

$$G(x, L) = 1 - \frac{x}{q^2} \int d^4 k \frac{k \cdot q}{k^2 G(x, \log k^2)(k+q)^2} - \dots \Big|_{q^2=\mu^2}$$


where $L = \log(q^2/\mu^2)$.

So

$$\gamma_1(x) = x - \gamma_1(x)(1 - 2x\partial_x)\gamma_1(x).$$


0-5

Vector field of $\gamma'_1(x)$




0-7

Solutions which die on the axis



$s = 1$



0-10

The $P(x) = x$ family

The last bastion of exact solutions,

$$\gamma_1(x) = x - \gamma_1(x)(1 - sx\partial_x)\gamma_1(x).$$

$$s = 1: \gamma_1(x) = x + xW(C \exp(-\frac{1+x}{x})),$$

$$s = 2: \exp\left(\frac{(1+\gamma_1(x))^2}{2x}\right) \sqrt{-x} + \text{erf}\left(\frac{1+\gamma_1(x)}{\sqrt{-2x}}\right) \frac{\sqrt{\pi}}{\sqrt{2}} = C$$


$$s = 3/2: A(X) - x^{1/3} 2^{1/3} A'(X) = C(B(X) - x^{1/3} 2^{1/3} B'(X)) \text{ where } X = \frac{1+\gamma_1(x)}{2^{2/3} x^{2/3}}$$

$$s = 3: (\gamma_1(x)+1)A(X) - 2^{2/3} A'(X) = C((\gamma_1(x)+1)B(X) - 2^{2/3} B'(X)) \text{ where } X = \frac{(1+\gamma_1(x))^2 + 2x}{2^{4/3} x^{2/3}}$$

where A is the Airy Ai function, B the Airy Bi function and W the Lambert W function.

0-9


$s = 3/2$



0-11

$s = 3$

Show s animation here.




0-12

0-13

$s = 1$ revisited


The solution $\gamma_1(x) = x$ appears to be a separatrix.



0-14

$s = 1$ revisited cont.

But then again, maybe not



0-15

QED as a single equation

By the Baker, Johnson, Willey analysis we can reduce to a single equation for the photon propagator.

$$2\gamma_1(x) = P(x) - \gamma_1(x)(1 - x\partial_x)\gamma_1(x)$$

$s = 1$ gives a term $B_+(\mathbb{I})$ independent of X to take into account the fact that the photon propagator can not be inserted into the one loop graph.

To 2 loops


$$P(x) = \frac{x}{3} + \frac{x^2}{4}$$

To 4 loops we need to correct the primitives for our setup

$$P(x) = \frac{x}{3} + \frac{x^2}{4} + (-0.0312 + 0.06037)x^3 + (-0.6755 + 0.05074)x^4$$


0-16

QED to 2 loops



0-17


Zoomed in



0-18


QED to 4 loops

At 4 loops $P(0.992\dots) = 0$ changing everything.




0-19

Zoomed in



0-20

We are here



0-21

References

- [1] D.J. Broadhurst and D. Kreimer. Exact solutions of Dyson-Schwinger equations for iterated one-loop integrals and propagator-coupling duality. *Nucl.Phys. B*, 600:403–422, 2001. arXiv:hep-th/0012146.
- [2] A. Connes and D. Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert problem. II: The beta-function, diffeomorphisms and the renormalization group. *Commun. Math. Phys.*, 216:215–241, 2001. arXiv:hep-th/0003188.

0-22