

Solutions to the Problems on Polynomials

- 1:** Let $p(x)$ be a polynomial over \mathbf{Z} with at least four distinct integral roots. Show that there is no integer k such that $p(k)$ is prime.

Solution: Suppose that $p(a) = p(b) = p(c) = p(d) = 0$ where a, b, c and d are distinct integers. Then we have $p(x) = (x-a)(x-b)(x-c)(x-d)h(x)$ for some polynomial $h(x)$. Thus for all $k \in \mathbf{Z}$, $p(k)$ is a multiple of each of the four distinct integers $(k-a)$, $(k-b)$, $(k-c)$ and $(k-d)$. Either one of these four integers is equal to zero, or two of these four integers are not equal to 0 or ± 1 . In either case, $p(k)$ is not prime.

- 2:** Let $p(x)$ be a polynomial over \mathbf{C} . Show that $p(x)$ is even if and only if there exists a polynomial $q(x)$ over \mathbf{C} such that $p(x) = q(x)q(-x)$.

Solution: Suppose that $p(x) = q(x)q(-x)$ for some polynomial q . Then we have $p(-x) = q(-x)q(x) = p(x)$, so p is even. Conversely, suppose that p is even. Let $p(x) = a_0 + a_1x + \cdots + a_nx^n$ where $a_n \neq 0$. Since p is even we have $p(x) = p(-x)$ for all x , so $a_0 + a_1x + a_2x^2 + \cdots + a_nx^n = a_0 - a_1x + a_2x^2 - \cdots + (-1)^n a_nx^n$ for all x . Comparing coefficients, we see that $0 = a_1 = a_3 = a_5 = \cdots$, so we have

$$p(x) = a_0 + a_2x^2 + a_4x^4 + \cdots + a_{2k}x^{2k} = r(x^2),$$

where $r(u) = a_0 + a_2u + a_4u^2 + \cdots + a_{2k}u^k$. Say $r(u) = a_{2k}(u-r_1)(u-r_2)\cdots(u-r_k)$. Then we have

$$p(x) = r(x^2) = a_{2k}(x^2-r_1)(x^2-r_2)\cdots(x^2-r_k) = (-1)^k a_{2k}(r_1-x^2)(r_2-x^2)\cdots(r_k-x^2).$$

Choose $b_i \in \mathbf{C}$ so that $b_0^2 = (-1)^k a_{2k}$ and $b_i^2 = r_i$ for $i \geq 1$. Then

$$p(x) = b_0^2(b_1^2-x^2)(b_2^2-x^2)\cdots(b_k^2-x^2) = q(-x)q(x),$$

where $q(x) = b_0(b_1+x)(b_2+x)\cdots(b_k+x)$.

- 3:** Let $p(x)$ be a polynomial over \mathbf{R} of odd degree. Show that $p(p(x))$ has at least as many real roots as $p(x)$.

Solution: Let a_1, a_2, \dots, a_n be the distinct real roots of $p(x)$. Note that since $p(x)$ has odd degree, it is onto, and so for each i , we can choose $x_i \in \mathbf{R}$ so that $p(x_i) = a_i$. Note that the numbers x_i are distinct (since $x_i = x_j \implies a_i = p(x_i) = p(x_j) = a_j$) and we have $p(p(x_i)) = p(a_i) = 0$ for all i .

- 4:** Let $p(x)$ be a monic polynomial over \mathbf{Z} with the property that there exist positive integers k and l such that none of the integers $p(k+i)$ with $i = 1, 2, \dots, l$ is divisible by l . Show that $p(x)$ has no rational roots.

Solution: Let a be a rational root of a monic polynomial p with integer coefficients, and let k and l be positive integers. Since p is monic, we have $a \in \mathbf{Z}$, so $p(x) = (x-a)q(x)$ for some monic polynomial $q(x)$ over \mathbf{Z} . For all i we have $p(k+i) = (k+i-a)q(a)$. Choose i with $1 \leq i \leq l$ so that $l \mid (k+i-a)$. Then we have $l \mid p(k+i)$.

- 5:** Let $p(x) = \sum_{k=0}^{2n} (-1)^k (2n+1-k)x^k$. Show that $p(x)$ has no real roots.

Solution: When $x \leq 0$, $p(x) = (2n+1) + (2n)|x| + (2n-1)|x|^2 + \cdots + 3|x|^{2n-2} + 2|x|^{2n-1} + |x|^{2n} \geq (2n+1) > 0$. Also, we have

$$p(x) = (2n+1) - (2n)x + (2n-1)x^2 - \cdots + 3x^{2n-2} - 2x^{2n-1} + x^{2n}, \text{ and}$$

$$xp(x) = (2n+1)x - (2n)x^2 + \cdots - 2x^{2n} + x^{2n+1}, \text{ so}$$

$$p(x)(1+x) = (2n+1) + x - x^2 + x^3 - \cdots + x^{2n+1} = (2n+1) + \frac{x(x^{2n+1}+1)}{x+1},$$

so that when $x > 0$ we have $p(x) > 0$. Thus $p(x) > 0$ for all x .

- 6:** Let $p(x)$ be a polynomial with non-negative real coefficients. Show that $p(a^2)p(b^2) \geq p(ab)^2$ for all $a, b \in \mathbf{R}$.

Solution: Let $p(x) = \sum_{i=0}^n c_i x^i$ with each $c_i \geq 0$. Let $u, v \in \mathbf{R}^{n+1}$ be the vectors with $u_i = \sqrt{c_i} a^i$ and $v_i = \sqrt{c_i} b^i$ for $0 \leq i \leq n$. Then $|u|^2 = u \cdot u = \sum_{i=0}^n c_i a^{2i} = p(a^2)$, and $|v|^2 = v \cdot v = \sum_{i=0}^n c_i b^{2i} = p(b^2)$, and $u \cdot v = \sum_{i=0}^n c_i a^i b^i = p(ab)$. By the Cauchy-Schwarz Inequality, $p(ab)^2 = (u \cdot v)^2 \leq |u|^2 |v|^2 = p(a^2)p(b^2)$.

7: Let $p(x)$ be a polynomial over \mathbf{Z} of degree at least 2. Show that there is a polynomial $q(x)$ over \mathbf{Z} such that $p(q(x))$ is reducible over \mathbf{Z} .

Solution: For $g(x) = p(x) - p(a)$ we have $g(a) = 0$ and so $(x - a) \mid g(x)$. Say $g(x) = (x - a)h(x)$, that is $p(x) - p(a) = (x - a)h(x)$. Note that since $\deg(p) \geq 2$ we have $\deg(h) \geq 1$. It follows that for any polynomial $f(x)$ we have $p(x) - p(f(x)) = (x - f(x))h(x)$. Take $f(x) = p(x) + x$ to get $p(x) - p(p(x) + x) = -p(x)h(x)$, so we have $p(p(x) + x) = p(x)(1 + h(x))$. Thus we can take $q(x) = p(x) + x$.

8: Let a and b be distinct real numbers. Solve $(z - a)^4 + (z - b)^4 = (a - b)^4$ for $z \in \mathbf{C}$.

Solution: We expand and simplify to get

$$z^4 - 4az^3 + 6a^2z^2 - 4a^3z + a^4 + z^4 - 4bz^3 + 6b^2z^2 - 4b^3z + b^4 = a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$$

$$2z^4 - 4(a+b)z^3 + 6(a^2+b^2)z^2 - 4(a^3+b^3)z + 4(a^3b+ab^3) - 6a^2b^2 = 0.$$

Thus we need to solve $f(z) = 0$ where $f(z) = z^4 - 2(a+b)z^3 + 3(a^2+b^2)z^2 - 2(a^3+b^3)z + (2a^3b - 3a^2b^2 + 2ab^3)$. Note that $f(a) = f(b) = 0$. Let a, b, c and d be the four roots of $f(z)$. Then we must have

$$a + b + c + d = 2(a + b) \text{ and}$$

$$abcd = ab(2a^2 - 3ab + 2b^2).$$

Thus we have $c + d = a + b$ and $cd = (2a^2 - 3ab + 2b^2)$, so c and d are the two solutions of the equation $z^2 - (a + b)z + (2a^2 - 3ab + 2b^2) = 0$, which are given by

$$c, d = \frac{(a + b) \pm \sqrt{(a^2 + 2ab + b^2) - 4(2a^2 - 3ab + 2b^2)}}{2}$$

$$= \frac{(a + b) \pm \sqrt{14ab - 7(a^2 + b^2)}}{2} = \frac{(a + b) \pm i(a - b)\sqrt{7}}{2}.$$

9: Let a_1, a_2, \dots, a_n be distinct integers. Show that $p(x) = \prod_{i=1}^n (x - a_i) - 1$ is irreducible.

Solution: Suppose, for a contradiction, that $p(x)$ is reducible. Say $p(x) = f(x)g(x)$ where f and g are both of degree less than n . For each i we have $p(a_i) = -1$ so $f(a_i)g(a_i) = -1$, and so either $f(a_i) = 1$ and $g(a_i) = -1$ or $f(a_i) = -1$ and $g(a_i) = 1$. Thus for all i we have we have $f(a_i) + g(a_i) = 0$. Since $f + g$ is of degree less than n , and the a_i are distinct, we must have $f(x) + g(x) = 0$ for all x . Since $g(x) = -f(x)$, we have $p(x) = f(x)g(x) = -f(x)^2$. But the coefficient of x^n in $p(x)$ is equal to 1, and the coefficient of x^n in $-f(x)^2$ is equal to -1 , so this is not possible.

10: Let $p_1(x) = x^2 - 2$ and for $k \geq 2$ let $p_k(x) = p_1(p_{k-1}(x))$. Show that the roots of $p_n(x) - x$ are real and distinct for all n .

Solution: Let $x(t) = 2 \cos t$ for $0 \leq t \leq \pi$. We have $p_1(x(t)) = (2 \cos t)^2 - 2 = 4 \cos^2 t - 2 = 2 \cos 2t$. Verify using mathematical induction that $p_n(x(t)) = 2 \cos(2^n t)$ for all $n \geq 1$. For $0 \leq t \leq \pi$ we have

$$p_n(x(t)) = x(t) \iff 2 \cos(2^n t) = 2 \cos t \iff \cos 2^n t = \cos t$$

$$\iff 2^n t = \pm t + 2\pi k, \text{ for some } k \in \mathbf{Z}$$

$$\iff t = \frac{2\pi k}{2^n \pm 1}, \text{ for some } k \in \{0, 1, 2, \dots, n - 1\}.$$

Thus we have found $2n$ distinct solutions $x(t)$ with $0 \leq t \leq \pi$, and since the polynomial $p_n(x) - x$ is of degree $2n$, these are all of the roots of $p_n(x) - x$.

11: Let $p(x) = \sum_{i=0}^n a_i x^i$ with $a_0 = a_n = 1$ and $a_i > 0$ for all i . Show that if $p(x)$ has n distinct real roots then $p(2) \geq 3^n$.

Solution: Since every $a_i \geq 0$, all of the roots of $p(x)$ must be negative. Say the roots are $-r_1, -r_2, \dots, -r_n$, so we have $p(x) = (x + r_1)(x + r_2) \cdots (x + r_n)$. Using the fact that $2 + r \geq 3r^{1/3}$ for all $r > 0$, and the fact that $r_1 r_2 \cdots r_n = p(0) = 1$, we have

$$p(2) = (2 + r_1)(2 + r_2) \cdots (2 + r_n) \geq (3r_1^{1/3})(3r_2^{1/3}) \cdots (3r_n^{1/3}) = 3^n (r_1 r_2 \cdots r_n)^{1/3} = 3^n.$$

12: Let $p(x)$ be the polynomial of degree n such that that $p(k) = \frac{k}{k+1}$ for $k = 0, 1, \dots, n$. Find $p(n+1)$.

Solution: Let $q(x) = (x+1)p(x) - x$. Then $\deg(q) = n+1$ and we have $q(k) = (k+1)\frac{k}{k+1} - k = 0$ for $0 \leq k \leq n$, and so $q(x) = c(x-0)(x-1)\cdots(x-n)$ for some constant c . Also, note that $q(-1) = -1$ so that $c(-1)(-2)\cdots(-1-n) = -1$, and so we have $c = \frac{(-1)^{n+1}}{(n+1)!}$. Thus $q(x) = \frac{(-1)^{n+1}}{(n+1)!}(x-0)(x-1)\cdots(x-n)$. In particular $q(n+1) = \frac{(-1)^{n+1}}{(n+1)!}(n+1)(n)\cdots(1) = (-1)^{n+1}$. Since $q(x) = (x+1)p(x) - x$, we have $p(x) = \frac{q(x)+x}{x+1}$, and in particular

$$p(n+1) = \frac{(-1)^{n+1} + (n+1)}{n+2} = \begin{cases} \frac{n}{n+2} & \text{if } n \text{ is even,} \\ 1 & \text{if } n \text{ is odd.} \end{cases}$$

13: Find all polynomials over \mathbf{C} whose coefficients are all equal to ± 1 and whose roots are all real.

Solution: Let $p(x) = \sum_{i=0}^n c_i x^i$ with each $c_i = \pm 1$, and suppose the roots of $p(x)$ are all real. Let the roots be a_1, a_2, \dots, a_n , repeated if necessary according to multiplicity, so that $p(x) = c_n(x-a_1)(x-a_2)\cdots(x-a_n)$. By partially expanding this product, we see that $c_0 = (-1)^n c_n \prod_{1 \leq i \leq n} a_i$, and $c_{n-1} = -c_n \sum_{1 \leq i \leq n} a_i$, and $c_{n-2} = c_n \sum_{1 \leq i < j \leq n} a_i a_j$. From the formulas for c_{n-1} and c_{n-2} we have

$$\sum_{1 \leq i \leq n} a_i^2 = \left(\sum_{1 \leq i < j \leq n} a_i a_j \right)^2 - 2 \sum_{1 \leq i \leq n} a_i a_j = \left(\frac{c_{n-1}}{c_n} \right)^2 - 2 \left(\frac{c_{n-2}}{c_n} \right).$$

Since each $c_i = \pm 1$ this gives $\sum_{1 \leq i \leq n} a_i^2 = 1 \pm 2$, and since $\sum_{1 \leq i \leq n} a_i^2 \geq 0$ we must have $\sum_{1 \leq i \leq n} a_i^2 = 1 + 2 = 3$.

By the Algebraic Geometric Mean Inequality, we have

$$1 = \sqrt[n]{1} = \sqrt[n]{c_0^2} = \sqrt[n]{\prod_{1 \leq i \leq n} a_i^2} \leq \frac{\sum_{1 \leq i \leq n} a_i^2}{n} = \frac{3}{n},$$

and so we must have $n \leq 3$. When $n = 1$ we find that all 4 polynomials $p(x) = \pm x \pm 1$ have real roots. When $n = 2$ we find that of the 8 polynomials $\pm x^2 \pm x \pm 1$, only the 4 polynomials $p(x) = \pm(x^2 \pm x - 1)$ have real roots. When $n = 3$, we must have equality in the Algebraic Geometric Mean Inequality, and this occurs when $a_1^2 = a_2^2 = a_3^2 = 1$. Of the 8 polynomials $\pm(x-1)^3, \pm(x-1)^2(x+1), \pm(x-1)(x+1)^2$ and $\pm(x+1)^3$, only the 4 polynomials $p(x) = \pm(x-1)^2(x+1), \pm(x-1)(x+1)^2$ have all coefficients ± 1 .

14: Find all polynomials $p(x)$ over \mathbf{R} such that $p(x)p(x+1) = p(x^2+x+1)$.

Solution: Let $p(x)$ be a polynomial over \mathbf{R} with $p(x)p(x+1) = p(x^2+x+1)$. If $p(x)$ is constant, say $p(x) = c$, then $c^2 = c$ so that $c = 0$ or 1 . Suppose that $p(x)$ is not constant. By replacing x by $x-1$, we see that $p(x-1)p(x) = p((x-1)^2+(x-1)+1) = p(x^2-x+1)$. Let $a \in \mathbf{C}$ be a root of $p(x)$ of largest possible norm. Then we have $p(a^2+a+1) = p(a)p(a+1) = 0$ and we have $p(a^2-a+1) = p(a-1)p(a) = 0$. Note that for any $0 \neq u \in \mathbf{C}$, one of the two complex numbers $u \pm a$ has larger norm than a (indeed, since $|u \pm a|^2 = |u|^2 \pm 2\operatorname{Re}(u\bar{a}) + |a|^2$, we see that if $\operatorname{Re}(u\bar{a}) \geq 0$ then $|u+a| > |a|$ and if $\operatorname{Re}(u\bar{a}) \leq 0$ then $|u-a| > |a|$). In particular, if $a \neq \pm i$ then $a^2+1 \neq 0$ and so one of $a^2 \pm a + 1$ has a larger norm than a , but since $a^2 \pm a + 1$ are both roots of $p(x)$, this would contradict our choice of a . Thus we must have $a = \pm i$. Say $p(x) = (x^2+1)^k q(x)$ where $q(\pm i) \neq 0$. Then since $(x^2+1)(x^2+2x+2) = (x^2+x+1)^2+1$ we have

$$q(x)q(x+1) = \frac{p(x)}{(x^2+1)^k} \cdot \frac{p(x+1)}{(x^2+2x+2)^k} = \frac{p(x^2+x+1)}{((x^2+x+1)^2+1)^k} = q(x^2+x+1)$$

and so $q(x)$ satisfies the same recursion as $p(x)$. As above, if $q(x)$ was not constant then its roots of largest norm would be $a = \pm i$, but we have $q(\pm i) \neq 0$ and so $q(x)$ must be constant, say $q(x) = c$. Also as above, we must have $q(x) = 0$ or $q(x) = 1$. But we cannot have $q(x) = 0$ since this would imply that $p(x) = (x^2+1)^k q(x) = 0$. Thus $q(x) = 1$ and $p(x) = (x^2+1)^k q(x) = (x^2+1)^k$.