

PMATH 347 Groups and Rings, Solutions to the Exercises for Chapter 2

1: In  $D_n$ , for  $k \in \mathbb{Z}_n$ , we write  $R_k$  for the rotation in about the point  $(0, 0)$  by the angle  $\frac{2\pi k}{n}$ , and we write  $F_k$  for the reflection in the line through  $(0, 0)$  and  $(\cos \frac{\pi k}{n}, \sin \frac{\pi k}{n})$ .

(a) Find all values of  $k \in \mathbb{Z}_6$  such that  $F_3 R_k F_1 = R_k$  in  $D_6$ .

Solution: We can use the formulas  $R_k R_l = R_{k+l}$ ,  $F_k F_l = R_{k-l}$ ,  $R_k F_l = F_{k+l}$  and  $F_k R_l = F_{k-l}$ . For  $k \in \mathbb{Z}_6$ ,  $F_3 R_k F_1 = R_k \iff F_{3-k} F_1 = R_k \iff R_{3-k-1} = R_k \iff 3-k-1 = k \iff 2k = 2 \iff k = 1$  or  $4$ .

(b) Find the centralizer of  $F_1$  in  $D_6$ .

Solution:  $R_k F_1 = F_1 R_k \iff F_{k+1} = F_{1-k} \iff k+1 = 1-k \iff 2k = 0 \iff k = 0$  or  $3$  in  $\mathbb{Z}_6$ . Also,  $F_k F_1 = F_1 F_k \iff R_{k-1} = R_{1-k} \iff k-1 = 1-k \iff 2k = 2 \iff k = 1$  or  $4$  in  $\mathbb{Z}_6$ . So  $C(F_1) = \{I, R_3, F_1, F_4\}$ .

2: (a) Find  $|GL(3, \mathbb{Z}_2)|$

Solution: For a matrix in  $GL(3, \mathbb{Z}_2)$ , the first row must be non-zero, and there are  $2^3 - 1 = 7$  such rows. Having fixed the first row, the second row can be any row that is not a multiple of the first; there are  $2^1$  multiples of the first row, so there are  $2^3 - 2^1 = 6$  possibilities for the second row. Having fixed the first two rows, the last row can be any row which is not a linear combination of the first two rows; there are  $2^2$  different linear combinations of the first two rows, so there are  $2^3 - 2^2 = 4$  possibilities for the last row. Altogether, there are  $7 \cdot 6 \cdot 4 = 168$  matrices in  $GL(3, \mathbb{Z}_2)$ , so  $|GL(3, \mathbb{Z}_2)| = 168$ .

(b) List all the elements in  $SO(3, \mathbb{Z}_2)$ .

Solution: Let  $A$  be the  $3 \times 3$  matrix over  $\mathbb{Z}_2$  with columns  $u_1, u_2, u_3$ . Note that  $A \in SO(3, \mathbb{Z}_2) = O(3, \mathbb{Z}_2) \iff A^T A = I \iff (u_k \cdot u_k = 1 \text{ for all } k \text{ and } u_k \cdot u_l = 0 \text{ for all } k \neq l)$ . The only vectors  $u_k$  with  $u_k \cdot u_k = 1$  are the three standard basis vectors  $e_k$  and the vector  $(1, 1, 1)^T$ , so each  $u_k$  must be one of these 4 vectors. Note that if  $u_k = (1, 1, 1)^T$  and  $u_l$  is any one of the above 4 vectors then  $u_k \cdot u_l = 1$ , not 0, so we cannot have  $u_k = (1, 1, 1)^T$ . Also, the vectors  $u_k$  must be distinct, so the 3 vectors  $u_k$  are equal to the 3 standard basis vectors (in some order). Thus there are 6 matrices in  $SO(3, \mathbb{Z}_2)$ , namely

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

3: (a) Show that  $U_{26}$  is cyclic.

Solution: Notice that  $\langle 7 \rangle = \{1, 7, 23, 5, 9, 11, 25, 19, 3, 21, 17, 15\} = U_{26}$ , and so  $U_{26}$  is cyclic.

(b) List all the elements and all the generators in every subgroup of  $U_{26}$ .

Solution: We list all the subgroups with the generators in boldface.

$$\begin{aligned} \langle 7 \rangle &= \{1, \mathbf{7}, 23, 5, 9, \mathbf{11}, 25, \mathbf{19}, 3, 21, 17, \mathbf{15}\} \\ \langle 7^2 \rangle &= \{1, \mathbf{23}, 9, 25, 3, \mathbf{17}\} \\ \langle 7^3 \rangle &= \{1, \mathbf{5}, 25, \mathbf{21}\} \\ \langle 7^4 \rangle &= \{1, \mathbf{9}, \mathbf{3}\} \\ \langle 7^6 \rangle &= \{1, \mathbf{25}\} \\ \langle 7^{12} \rangle &= \{\mathbf{1}\} \end{aligned}$$

4: (a) Determine the number of subgroups of  $\mathbb{Z}_{12,000}$ .

Solution: Since  $12,000 = 2^5 3^1 5^3$ , the divisors of 12,000 are of the form  $2^i 3^j 5^k$  with  $0 \leq i \leq 5$ ,  $0 \leq j \leq 1$  and  $0 \leq k \leq 3$ . Since there are 6 possible values for  $i$ , 2 possible values for  $j$  and 4 possible values for  $k$ , there are  $6 \cdot 2 \cdot 4 = 48$  divisors of 12,000. Thus there are 48 subgroups of  $\mathbb{Z}_{12,000}$ .

(b) Find the number of elements of even order in  $\mathbb{Z}_{12,000}$ .

Solution: The odd factors of 12,000 are of the form  $3^j 5^k$  with  $0 \leq j \leq 1$  and  $0 \leq k \leq 3$ . There are 8 such odd factors, namely 1, 5, 25, 125, 3, 15, 75 and 375, and correspondingly there are 8 subgroups of odd order in  $\mathbb{Z}_{12,000}$ . The elements in  $\mathbb{Z}_{12,000}$  of odd order are the generators of these 8 subgroups, so the number of elements of odd order is  $\phi(1) + \phi(5) + \phi(25) + \phi(125) + \phi(3) + \phi(15) + \phi(75) + \phi(375) = 1 + 4 + 20 + 100 + 2 + 8 + 40 + 200 = 375$ . Thus the number of elements of even order in  $\mathbb{Z}_{12,000}$  is  $12,000 - 375 = 825$ .

5: (a) Find the number of elements of each order in  $\mathbb{Z}_3 \times \mathbb{Z}_6$ .

Solution: There is 1 element of order 1, 1 of order 2, 8 of order 3 and 8 of order 6.

(b) List all the elements in every cyclic subgroup of  $\mathbb{Z}_3 \times \mathbb{Z}_6$ .

Solution: By the result of Part (a), there is 1 cyclic subgroup of order 1, 1 of order 2, 4 of order 3 and 4 of order 6. The cyclic subgroups are

$$\begin{aligned}\langle(0, 0)\rangle &= \{(0, 0)\} \\ \langle(0, 3)\rangle &= \{(0, 0), (0, 3)\} \\ \langle(0, 2)\rangle &= \{(0, 0), (0, 2), (0, 4)\} \\ \langle(1, 0)\rangle &= \{(0, 0), (1, 0), (2, 0)\} \\ \langle(1, 2)\rangle &= \{(0, 0), (1, 2), (2, 4)\} \\ \langle(1, 4)\rangle &= \{(0, 0), (1, 4), (2, 2)\} \\ \langle(0, 1)\rangle &= \{(0, 0), (0, 1), (0, 2), (0, 3), (0, 4), (0, 5)\} \\ \langle(1, 3)\rangle &= \{(0, 0), (1, 3), (2, 0), (0, 3), (1, 0), (2, 3)\} \\ \langle(1, 1)\rangle &= \{(0, 0), (1, 1), (2, 2), (0, 3), (1, 4), (2, 5)\} \\ \langle(1, 5)\rangle &= \{(0, 0), (1, 5), (2, 4), (0, 3), (1, 2), (5, 1)\}\end{aligned}$$

(c) List all the elements in every non-cyclic subgroup of  $\mathbb{Z}_3 \times \mathbb{Z}_6$ . Explain why your list is complete.

Solution:  $\mathbb{Z}_3 \times \mathbb{Z}_6$  and the subgroup  $\mathbb{Z}_3 \times \langle 2 \rangle = \{(0, 0), (1, 0), (2, 0), (0, 2), (1, 2), (2, 2), (0, 4), (1, 4), (2, 4)\}$  are non-cyclic subgroups of  $\mathbb{Z}_3 \times \mathbb{Z}_6$ . We now show that these are the only two. Since  $|\mathbb{Z}_3 \times \mathbb{Z}_6| = 18$ , any subgroup must have order 1, 2, 3, 6, 9 or 18. Any group of order 1, 2 or 3 must be cyclic, and any abelian group of order 6 is cyclic, so any non-cyclic subgroup of  $\mathbb{Z}_3 \times \mathbb{Z}_6$  must have order 9 or 18. Of course, the only subgroup of  $\mathbb{Z}_3 \times \mathbb{Z}_6$  of order 18 is  $\mathbb{Z}_3 \times \mathbb{Z}_6$  itself. Now, let  $H$  be a subgroup of order 9 in  $\mathbb{Z}_3 \times \mathbb{Z}_6$ . Then the elements of  $H$  could only be of order 1, 3 or 9. No elements in  $\mathbb{Z}_3 \times \mathbb{Z}_6$  have order 9, so  $H$  must consist of the identity along with 8 elements of order 3. But the group  $\mathbb{Z}_3 \times \mathbb{Z}_6$  only has 8 elements of order 3, so all of these must be in  $H$ , and hence  $H$  must be the group  $\mathbb{Z}_3 \times \langle 2 \rangle$ .