

PMATH 347 Groups and Rings, Solutions to the Exercises for Chapter 5

1: (a) List all the elements in every coset of $H = \{1, 9, 17, 33\}$ in $G = U_{40}$.

Solution: Since $|H| = 4$ and $|G| = 16$, we know there must $|G|/|H| = 4$ distinct cosets. They are $1H = \{1, 9, 17, 33\}$, $3H = \{3, 27, 11, 19\}$, $39H = (-1)H = \{-1, -9, -17, -33\} = \{39, 31, 23, 7\}$ and $37H = (-3)H = \{-3, -27, -11, -19\} = \{37, 13, 29, 21\}$.

(b) List all the elements in every left coset and every right coset of $H = \langle(1234)\rangle$ in $G = S_4$.

Solution: $G = \{(1), (12), (13), (14), (23), (24), (34), (12)(34), (13)(24), (14)(23), (123), (124), (132), (134), (142), (143), (234), (243), (1234), (1243), (1324), (1342), (1423), (1432)\}$ and $H = \{(1), (1234), (13)(24), (1432)\}$, and there are $24/4=6$ left cosets;

$$\begin{aligned}(1)H &= \{(1), (1234), (13)(24), (1432)\} \\ (12)H &= \{(12), (234), (1324), (143)\} \\ (13)H &= \{(13), (12)(34), (24), (14)(23)\} \\ (14)H &= \{(14), (123), (1342), (243)\} \\ (23)H &= \{(23), (134), (1243), (142)\} \\ (34)H &= \{(34), (124), (1423), (132)\}\end{aligned}$$

There are also 6 right cosets:

$$\begin{aligned}H(1) &= \{(1), (1234), (13)(24), (1432)\} \\ H(12) &= \{(12), (134), (1423), (243)\} \\ H(13) &= \{(13), (14)(23), (24), (12)(34)\} \\ H(14) &= \{(14), (234), (1243), (132)\} \\ H(23) &= \{(23), (124), (1342), (143)\} \\ H(34) &= \{(34), (123), (1324), (142)\}\end{aligned}$$

2: Find four distinct subgroups $G_i \leq S_5$, with $i = 1, 2, 3, 4$, such that $\text{orb}_{G_i}(1) = \{1, 2, 4\}$. For each of the four subgroups G_i , find $\text{stab}_{G_i}(1)$.

Solution: The groups G_i will permute the numbers 1, 2 and 4 amongst one another, and might also permute the numbers 3 and 5 amongst each other. The four groups are

$$\begin{aligned}G_1 &= \{(1), (124), (142)\} \\ G_2 &= \{(1), (124), (142), (12), (14), (24)\} \\ G_3 &= \{(1), (124), (142), (35), (124)(35), (142)(35)\} \\ G_4 &= \{(1), (124), (142), (12), (14), (24), (35), (124)(35), (142)(35), (12)(35), (14)(35), (24)(35)\}\end{aligned}$$

and the stabilizers are the groups $\text{stab}_{G_1}(1) = \{(1)\}$, $\text{stab}_{G_2}(1) = \{(1), (24)\}$, $\text{stab}_{G_3}(1) = \{(1), (35)\}$ and $\text{stab}(1) = \{(1), (24), (35), (24)(35)\}$.

3: Let $G = \mathbb{Z}^2$ and let $H = \langle (3, 2), (6, 8) \rangle = \{k(3, 2) + l(6, 8) \mid k, l \in \mathbb{Z}\}$. For each pair $(a, b) \in G$ with $0 \leq a < 6$ and $0 \leq b < 2$, find the order of $(a, b) + H$ in the group G/H .

Solution: The order of $(a, b) + H$ in G/H is the smallest positive integer n such that $n(a, b) \in H$, so we would like to find an easy way to test a given point (c, d) to determine whether it is in H . One way is to use a picture of the group H in G to determine whether a given point $(c, d) \in H$. Another is as follows.

$$\begin{aligned} (c, d) \in H &\iff (c, d) = k(3, 2) + l(6, 8) \quad \text{for some } (k, l) \in \mathbb{Z}^2 \\ &\iff \begin{pmatrix} 3 & 6 \\ 2 & 8 \end{pmatrix} \begin{pmatrix} k \\ l \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix} \quad \text{for some } (k, l) \in \mathbb{Z}^2 \\ &\iff \begin{pmatrix} k \\ l \end{pmatrix} = \begin{pmatrix} 3 & 6 \\ 2 & 8 \end{pmatrix}^{-1} \begin{pmatrix} c \\ d \end{pmatrix} = \frac{1}{12} \begin{pmatrix} 8 & -6 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} \in \mathbb{Z}^2 \\ &\iff 8c - 6d = 0 \pmod{12} \quad \text{and} \quad -2c + 3d = 0 \pmod{12} \\ &\iff 2c = 3d \pmod{12} \end{aligned}$$

For $(a, b) \in \mathbb{Z}^2$ let us write $[a, b] = (a, b) + H \in G/H$. With the help of a picture of H , or by using the formula $(c, d) \in H \iff 2c = 3d \pmod{12}$, we find that $|[1, 1]| = 12$ with $\langle [1, 1] \rangle = \{[0, 0], [1, 1], [2, 2], \dots, [11, 11]\}$ (note that $[12, 12] = [0, 0]$ in G/H since $(12, 12) \in H$). Consequently we have $|[k, k]| = 12/\gcd(k, 12)$ for all k . We could find the order of $[a, b]$ for each of the given elements (a, b) in a similar manner, but it is more amusing to notice that every element $[a, b]$ is equal to $[k, k]$ for some $0 \leq k < 12$. For example, we have $[1, 0] = [1, 0] + [9, 10] = [10, 10]$ (since $[9, 10] \in H$), and so $|[1, 0]| = |[10, 10]| = \frac{12}{\gcd(10, 12)} = 6$. In this way, we can find the order of each of the given elements: $|[0, 0]| = 1$, $|[1, 0]| = |[10, 10]| = 6$, $|[2, 0]| = |[8, 8]| = 3$, $|[3, 0]| = |[6, 6]| = 2$, $|[4, 0]| = |[4, 4]| = 3$, $|[5, 0]| = |[2, 2]| = 6$, $|[0, 1]| = |[3, 3]| = 4$, $|[1, 1]| = 12$, $|[2, 1]| = |[11, 11]| = 12$, $|[3, 1]| = |[9, 9]| = 4$, $|[4, 1]| = |[7, 7]| = 12$ and $|[5, 1]| = |[5, 5]| = 12$

4: Determine which of the following five subgroups of S_4 are normal: $\langle (12) \rangle$, $\langle (12)(34) \rangle$, $\langle (123) \rangle$, $\langle (1234) \rangle$ and $\text{stab}_{S_4}(1)$.

Solution: Notice that we have $(14)(12)(14) = (24) \notin \langle (12) \rangle$, $(14)(12)(34)(14) = (13)(24) \notin \langle (12)(34) \rangle$, $(14)(123)(14) = (234) \notin \langle (123) \rangle$, $(14)(1234)(14) = (1423) \notin \langle (1234) \rangle$, and $(14)(24)(14) = (12) \notin \text{stab}_{S_4}(1)$, and so none of these subgroups are normal.