

## §1.6 - Compositions and Inverses

Given two functions  $f(x)$  and  $g(x)$ , the composition of  $f$  with  $g$ , denoted  $f \circ g$ , is the function

$$(f \circ g)(x) = f(g(x))$$

Similarly,

$$(g \circ f)(x) = g(f(x))$$

Ex: Let  $f(x) = x^2 + 1$  and  $g(x) = x + 7$ .

$$(f \circ g)(1) = f(g(1)) = f(1+7) = f(8) = 8^2 + 1 = \boxed{65}$$

$$(g \circ f)(1) = g(f(1)) = g(1^2 + 1) = g(2) = 2 + 7 = \boxed{9}$$

$$(f \circ g)(x) = f(g(x)) = f(x+7) = \boxed{(x+7)^2 + 1}$$

$$(g \circ f)(x) = g(f(x)) = g(x^2 + 1) = (x^2 + 1) + 7 = \boxed{x^2 + 8}$$

Note: We can also talk about  $f \circ f$ ,  $g \circ g$ ,  $g \circ f \circ g$ , etc.

The domain of  $f(g(x))$  is found as before (i.e., by removing any "problem points"), but now we must also account for the domain of  $g(x)$ , the inner function, by removing its "problem points".

Ex: Let  $f(x) = \frac{x}{1+x}$ ,  $g(x) = \frac{1}{x}$ . Find  $f(g(x))$  and its domain.

Solution:  $f(g(x)) = f\left(\frac{1}{x}\right) = \frac{\frac{1}{x}}{1 + \frac{1}{x}} = \frac{\frac{1}{x}}{\frac{x}{x} + \frac{1}{x}} = \frac{\frac{1}{x}}{\frac{x+1}{x}}$

Must exclude  $x=0$  since it's not in the domain of  $g$ !

Must exclude  $x=-1$ , else we divide by 0

$$= \frac{1}{\cancel{x}} \cdot \frac{\cancel{x}}{x+1}$$

$$= \frac{1}{x+1}$$

$$\therefore \text{Domain} = \{x \in \mathbb{R} : x \neq 0 \text{ and } x \neq -1\}$$

Ex: If  $f(x) = \sqrt{x} + 1$  and  $g(x) = (x-1)^2 + 1$ , find  $g \circ f$  as well as its domain and range.

Solution:  $g(f(x)) = g(\sqrt{x} + 1) = ((\sqrt{x} + 1) - 1)^2 + 1$   
 $= (\sqrt{x})^2 + 1 = \boxed{x + 1}$

Domain: All  $x \in \mathbb{R}$ , but we also need  $x \geq 0$  for the domain of  $f$ , hence  $\boxed{\text{domain} = [0, \infty)}$ .

Range: Since  $y = g(f(x)) = x + 1$  with  $x \in [0, \infty)$ ,  
 $\boxed{\text{range} = [1, \infty)}$ .

## Inverse Functions

Given an input  $x$ , a function  $y = f(x)$  tells us the corresponding  $y$ -value. But what if we start with  $y$  and want the corresponding  $x$ ?

Ex:  $y = f(x) = \frac{x+3}{x}$ . Find  $x$  such that  $y=7$

Solution:  $y = 7 = \frac{x+3}{x} \Rightarrow 7x = x+3$   
 $\Rightarrow 6x = 3$   
 $\Rightarrow x = \frac{3}{6} = \boxed{\frac{1}{2}}$

In fact, we can do this with any  $y$ :

$$y = \frac{x+3}{x} \Rightarrow xy = x+3$$
$$\Rightarrow xy - x = 3$$
$$\Rightarrow x(y-1) = 3 \Rightarrow \boxed{x = \frac{3}{y-1}}$$

This new function is called the inverse of  $f$  and

is written  $x = f^{-1}(y)$ . It "undoes" the function  $f$ !

e.g. With  $y = f(x) = \frac{x+3}{x}$ , if  $y=7$ , then

$$x = f^{-1}(y) = \frac{3}{y-1} = \frac{3}{7-1} = \boxed{\frac{1}{2}}$$

(same as before!)

Note: We often like seeing  $x$  as the input variable and  $y$  as the output variable, even for inverses.

So, we usually swap  $x$  and  $y$  when calculating  $f^{-1}$ .

e.g. In our previous example, we have

$$f(x) = \frac{x+3}{x} \Rightarrow f^{-1}(x) = \frac{3}{x-1} \quad \leftarrow \begin{array}{l} f^{-1}(y) = \frac{3}{y-1} \\ \text{swap!} \end{array}$$

Ex: Find  $f^{-1}(x)$  given  $f(x) = \frac{2x}{3x-1}$ .

Solution:

$$\begin{aligned} y &= \frac{2x}{3x-1} \Rightarrow y(3x-1) = 2x \\ &\Rightarrow 3xy - y = 2x \\ &\Rightarrow x(3y-2) = y \\ &\Rightarrow x = \frac{y}{3y-2} \end{aligned}$$

Swapping the variables:  $y = f^{-1}(x) = \frac{x}{3x-2}$

Q: Do all functions have inverses? No!


Ex: Consider  $y = f(x) = x^2$ .

There is no way to "undo"

$f$  since  $y=4$ , for example,



could have come from

multiple  $x$  values:  $x=2$  or  $x=-2$ .




For a function  $y = f(x)$  to have an inverse, it must pass the horizontal line test: every horizontal line intersects the graph of  $f$  at most once.

To get an inverse for  $f(x) = x^2$ , we would need to restrict its domain



Graphically,  $f^{-1}(x)$  can be obtained by reflecting the graph of  $f(x)$  over the line  $y=x$ .



- Useful facts:
- Domain of  $f$  = Range of  $f^{-1}$
  - Range of  $f$  = Domain of  $f^{-1}$ .

Ex: What is the range of  $f(x) = \frac{2x}{3x-1}$  ?

Solution: We showed earlier that  $f^{-1}(x) = \frac{x}{3x-2}$

The domain of  $f^{-1}$  is  $\{x \in \mathbb{R} : x \neq \frac{2}{3}\}$  and

hence the range of  $f$  is  $\{y \in \mathbb{R} : y \neq \frac{2}{3}\}$ .



